arXiv:1110.0401v2 [math.PR] 2 Mar 2012

FIRST PASSAGE TIMES FOR SUBORDINATE BROWNIAN MOTIONS

MATEUSZ KWAŚNICKI, JACEK MAŁECKI, MICHAŁ RYZNAR

Abstract. Let X_t be a subordinate Brownian motion, and suppose that the Lévy measure of the underlying subordinator has a completely monotone density. Under very mild conditions, we find integral formulae for the tail distribution $\mathbf{P}(\tau_x > t)$ of first passage times τ_x through a barrier at $x > 0$, and its derivatives in t. As a corollary, we examine the asymptotic behaviour of $\mathbf{P}(\tau_x > t)$ and its t-derivatives, either as $t \to \infty$ or $x \to 0^+$.

1. Introduction

The present article complements and extends the results of the recent paper [28], where spectral theory for a class of Lévy processes killed upon leaving a half-line was developed. In a closely related paper [29], first passage times were studied for a rather general class of one-dimensional Lévy processes. In the present article, more detailed properties of first passage times are established for processes considered in [28]: symmetric Lévy processes, whose Lévy measure has a completely monotone density function on $(0, \infty)$. More precisely, we prove asymptotic fomulae, regularity, and estimates of the tail distribution $\mathbf{P}(\tau_x > t)$ of the first passage time through a barrier at the level x for a Lévy process X_t:

$$\tau_x = \inf \{t \ge 0 : X_t \ge x\}, \qquad\qquad x \ge 0,$$

as well as its derivatives in t. Alternatively, the results can be stated in terms of the *supremum functional* $M_t = \sup_{s \in [0,t]} X_s$, since we have $\mathbf{P}(\tau_x > t) = \mathbf{P}(M_t < x)$ for all $t, x \ge 0$.

In [28], a formula was given for generalised eigenfunctions $F_\lambda(x)$ of the transition semigroup of the killed process. As an application, the distribution of first passage times was expressed in terms of the eigenfunctions $F_\lambda(x)$. The full statement of this result was only announced, and a formal proof was given under more restrictive conditions. In the present paper, we provide the proof in the general case (Theorem 1.6). The expression for the distribution of τ_x is then used to find estimates and asymptotic expansion of $(d/dt)^n \mathbf{P}(\tau_x > t)$. This requires detailed analysis of the eigenfunctions $F_\lambda(x)$.

The double Laplace transform (in t and x) of $\mathbf{P}(\tau_x > t)$ is known for general Lévy processes since 1957 due to the result of Baxter and Donsker (Theorem 1 in [4]). For symmetric Lévy processes X_t with Lévy-Khintchin exponent $\Psi(\xi)$,

$$\begin{aligned} &\int_0^\infty \int_0^\infty e^{-\xi x - zt} \mathbf{P}(\tau_x > t) dx dt \\ &\qquad = \frac{1}{\xi \sqrt{z}} \exp\left(-\frac{1}{\pi} \int_0^\infty \frac{\xi \, \log(z + \Psi(\zeta))}{\xi^2 + \zeta^2} \, d\zeta \right). \end{aligned} \tag{1.1}$$

However, the double Laplace transform in (1.1) has been inverted only for few special cases. It is a classical result that for the Brownian motion, τ_x is the (1/2)-stable subordinator. An explicit formula for the distribution of τ_x was found for the Cauchy process (the symmetric 1-stable process) by Darling [10], for a compound Poisson process with

---

Mateusz Kwaśnicki received financial support of the Foundation for Polish Science.

$\Psi(\xi) = 1 - \cos\xi$ by Baxter and Donsker [4], and for the Poisson process with drift by Pyke [32]. A formula for the single Laplace transform for symmetric Lévy processes, under some mild assumptions, was given recently in [29] (see Theorem 1.3 below). In the development of the fluctuation theory for Lévy processes, many new identities involving first passage times were derived (see [6, 12, 30, 33] for a general account on fluctuation theory), including various other characterisations of $\mathbf{P}(\tau_x > t)$, at least in the stable case, see [5, 7, 10, 11, 14, 15, 16, 18, 19, 26, 27, 36]. First passage times τ_x and the supremum functional M_t play an important role in many areas of applied probability ([2, 3]), mathematical physics ([20, 25]), and also in potential theory of Lévy processes ([9, 21, 22, 23, 24]).

The main result of this article is an explicit, applicable for numerical computations expression for $\mathbf{P}(\tau_x > t)$ for a class of symmetric Lévy processes, which includes symmetric α-stable processes, relativistic α-stable processes and geometric α-stable processes (in particular, the variance gamma process) and many others. More precisely, the following assumption is in force throughout the article.

Assumption 1.1. Any of the following equivalent conditions is satisfied (see Proposition 2.13 in [28]):

(a) X_t is a *subordinate Brownian motion*, $X_t = B_{Z_t}$, and the Lévy measure of the subordinator Z_t has a completely monotone density. Here B_s is the one-dimensional Brownian motion ($\operatorname{Var} B_s = 2s$), Z_t is a *subordinator* (nonnegative Lévy process), and B_s and Z_t are independent processes;
(b) X_t is a symmetric Lévy process, whose Lévy measure has a completely monotone density on $(0, \infty)$;
(c) X_t is a Lévy process with Lévy-Khintchine exponent $\Psi(\xi) = \psi(\xi^2)$ for some *complete Bernstein function* $\psi(\xi)$.

We assume that X_t is non-trivial, that is, X_t is not constantly 0.

Remark 1.2. All explicit formulae and estimates proved in this article are given in terms of the complete Bernstein function $\psi(\xi)$. Translation to the Lévy-Khintchine exponent $\Psi(\xi) = \psi(\xi^2)$ is immediate, but usually results in less elegant expressions.

In this article, the term *explicit formula* is used for an expression involving a finite number of (absolutely convergent) integrals, elementary functions and the function ψ. The complete Bernstein function ψ extends to a holomorphic function on $\mathbf{C} \setminus (-\infty, 0]$ (see Preliminaries). Sometimes (namely, in the formula for $F_\lambda(x)$) we also use this holomorphic extension of ψ. □

Our proofs are based on the following two theorems.

Theorem 1.3 (Corollary 4.2 in [29]). *We have*

$$\begin{aligned}\int_0^\infty e^{-\xi x} \mathbf{P}(\tau_x > t) dx = \frac{2}{\pi} \int_0^\infty & \frac{\lambda}{\lambda^2 + \xi^2} \, \frac{\psi'(\lambda^2)}{\sqrt{\psi(\lambda^2)}} \times \\ & \times \exp\left(\frac{1}{\pi} \int_0^\infty \frac{\xi \log \frac{\lambda^2 - \zeta^2}{\psi(\lambda^2) - \psi(\zeta^2)}}{\xi^2 + \zeta^2} \, d\zeta \right) e^{-t\psi(\lambda^2)} d\lambda\end{aligned} \tag{1.2}$$

for all $t, \xi > 0$. □

We remark that the above result is proved in [29] for all symmetric Lévy processes with Lévy-Khintchine exponent $\Psi(\xi)$ having strictly positive derivative on $(0, \infty)$.

The transition semigroup $P_t^{(0,\infty)}$ (acting on $L^p((0, \infty))$ for any $p \in [1, \infty]$) of the process X_t killed upon leaving the half-line $(0, \infty)$, and its $L^2((0, \infty))$ generator $\mathcal{A}_{(0,\infty)}$, are defined formally, for example, in [28]. These notions are only required in the statements of Theorems 1.4 and 1.10, and therefore they are not discussed in detail below.

Theorem 1.4 (Theorem 1.1 in [28])**.** *For every $\lambda > 0$, there is a bounded continuous function F_λ on $(0, \infty)$ which is the eigenfunction of $P_t^{(0,\infty)}$, that is,*

$$P_t^{(0,\infty)} F_\lambda(x) = e^{-t\psi(\lambda^2)} F_\lambda(x)$$

for all $t, x > 0$. The function F_λ is characterised by its Laplace transform:

$$\mathcal{L}F_\lambda(\xi) = \frac{\lambda}{\lambda^2 + \xi^2} \exp\left(\frac{1}{\pi}\int_0^\infty \frac{\xi}{\xi^2+\zeta^2} \log\frac{\psi'(\lambda^2)(\lambda^2-\zeta^2)}{\psi(\lambda^2)-\psi(\zeta^2)}\,d\zeta\right) \tag{1.3}$$

for $\xi \in \mathbf{C}$ such that $\operatorname{Re}\xi > 0$*. Furthermore, for $x > 0$ we have*

$$F_\lambda(x) = \sin(\lambda x + \vartheta_\lambda) - G_\lambda(x), \tag{1.4}$$

where the phase shift ϑ_λ *belongs to* $[0, \pi/2)$*, and the* correction term $G_\lambda(x)$ *is a bounded, completely monotone function on $(0,\infty)$. More precisely, we have*

$$\vartheta_\lambda = -\frac{1}{\pi}\int_0^\infty \frac{\lambda}{\lambda^2-\zeta^2} \log\frac{\psi'(\lambda^2)(\lambda^2-\zeta^2)}{\psi(\lambda^2)-\psi(\zeta^2)}\,d\zeta, \tag{1.5}$$

and G_λ is the Laplace transform of a finite measure γ_λ on $(0,\infty)$. When $\psi(\xi)$ extends to a function $\psi^+(\xi)$ holomorphic in the upper complex half-plane $\{\xi \in \mathbf{C} : \operatorname{Im}\xi > 0\}$ and continuous in $\{\xi \in \mathbf{C} : \operatorname{Im}\xi \ge 0$, and furthermore $\psi^+(-\xi) \ne \psi(\lambda)$ for all $\xi > 0$, then the measure γ_λ is absolutely continuous, and

$$\begin{aligned}\gamma_\lambda(d\xi) = \frac{1}{\pi}&\left(\operatorname{Im}\frac{\lambda\psi'(\lambda^2)}{\psi(\lambda^2)-\psi^+(-\xi^2)}\right)\\ &\times \exp\left(-\frac{1}{\pi}\int_0^\infty \frac{\xi}{\xi^2+\zeta^2}\log\frac{\psi'(\lambda^2)(\lambda^2-\zeta^2)}{\psi(\lambda^2)-\psi(\zeta^2)}\,d\zeta\right) d\xi\end{aligned} \tag{1.6}$$

for $\xi > 0$. □

We introduce the following two conditions:

$$\sup_{\xi>0}\frac{\xi|\psi''(\xi)|}{\psi'(\xi)} < 2, \tag{1.7}$$

and, given $t_0 > 0$,

$$\int_1^\infty \sqrt{\frac{\psi'(\xi^2)}{\psi(\xi^2)}}\, e^{-t_0\psi(\xi^2)} d\xi < \infty. \tag{1.8}$$

Remark 1.5. (a) By Proposition 2.3(b), for every complete Bernstein function ψ the supremum in (1.7) is not greater than 2. Condition (1.7), is needed only to assert that $\sup_{\lambda>0}\vartheta_\lambda < \pi/2$, and can be replaced by the latter.

(b) When $\psi(\xi)$ is *unbounded* (that is, X_t is not a compound Poisson process) and *regularly varying* of order ϱ_0 at 0 and ϱ_∞ at ∞, then $\varrho_0, \varrho_\infty \in [0,1]$ and $\lim_{\xi\to 0^+}(\xi|\psi''(\xi)|/\psi'(\xi)) = 1 - \varrho_0$, $\lim_{\xi\to\infty}(\xi|\psi''(\xi)|/\psi'(\xi)) = 1 - \varrho_\infty$ (see [8]). Hence (1.7) is automatically satisfied.

(c) Assumption (1.8) is a rather mild growth condition on $\psi(\xi)$ for large ξ. By Proposition 2.3(a), $\sqrt{\psi'(\xi^2)/\psi(\xi^2)} \le 1/\xi$. Hence, (1.8) is satisfied for all $n \ge 0$ an all $t_0 > 0$ whenever $\psi(\xi) \ge c\log(\xi)$ ($\xi > 0$) for some $c > 0$. When $\psi(\xi) = f(\log(1+\xi))$, then (1.8) is equivalent to $\int_1^\infty \sqrt{f'(s)/f(s)}\, e^{-t_0 f(s)} ds < \infty$. For example, if $\psi(\xi) = \log(1+\log(1+\xi))$, then (1.8) holds if and only if $t_0 > 1/2$. □

In particular, many processes frequently found in literature, including symmetric α-stable processes, relativistic α-stable processes and geometric α-stable processes, satisfy (1.7) and (1.8). These and some other examples are discussed in Section 7.

The following are the main results of the article. The first of them provides a formula for $\mathbf{P}(\tau_x > t)$ and its derivatives in t, and it was proved in [28] under more restrictive assumptions. The full statement of the theorem (for $n = 0$ and $n = 1$) was announced in [28] as Theorem 1.8.

Theorem 1.6. *If* (1.7) *and* (1.8) *hold for some* $t_0 > 0$*, then for all* $n \ge 0$*,* $t > t_0$ *(*$t \ge t_0$ *if* $n = 0$*) and* $x > 0$*,*

$$\begin{aligned}(-1)^n \frac{d^n}{dt^n} \mathbf{P}(\tau_x > t) \\ = \frac{2}{\pi} \int_0^\infty \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}} \, (\psi(\lambda^2))^n e^{-t\psi(\lambda^2)} F_\lambda(x) d\lambda.\end{aligned} \tag{1.9}$$

In Theorem 4.6 in [29] it is proved that

$$\frac{1}{20000} \min\left(1, \frac{1}{\sqrt{t\psi(1/x^2)}}\right) \le \mathbf{P}(\tau_x > t) \le 10 \min\left(1, \frac{1}{\sqrt{t\psi(1/x^2)}}\right) \tag{1.10}$$

for *all* $t, x > 0$ for a relatively wide class of symmetric Lévy processes X_t (similar results for many asymmetric processes are also available in [29]). Our second result generalises these estimates to derivatives of $\mathbf{P}(\tau_x > t)$ in t, for a restricted class of processes and t large enough. For similar bounds with slightly different assumptions, see Lemma 6.2, Corollary 6.4, Remark 6.5 and Proposition 7.1.

Theorem 1.7. (a) *If* (1.7) *and* (1.8) *hold for some* $t_0 > 0$*, then the distribution of* τ_x *is* ultimately completely monotone*, that is, for each fixed* $n \ge 0$*,* $(-1)^n (d^n/dt^n) \mathbf{P}(\tau_x > t) \ge 0$ *for* t *large enough.*

(b) *If* $\varrho = \sup_{\xi > 0} (\xi |\psi''(\xi)| / \psi'(\xi)) < 1$ *(cf.* (1.7)*), then there are positive constants* $c_1(n)$*,* $c_2(n)$*,* $c_3(n, \varrho)$ *such that*

$$\frac{c_1(n)}{t^{n+1/2} \sqrt{\psi(1/x^2)}} \le (-1)^n \frac{d^n}{dt^n} \mathbf{P}(\tau_x > t) \le \frac{c_2(n)}{t^{n+1/2} \sqrt{\psi(1/x^2)}} \tag{1.11}$$

for $n \ge 0$*,* $x > 0$ *and* $t \ge c_3(n, \varrho)/\psi(1/x^2)$*. When* $c_2(n)$ *is replaced by a constant* $\tilde{c}_2(n, \varrho)$*, then the upper bound in* (1.11) *holds for all* $t, x > 0$*.*

Next, we study the asymptotic behaviour of $\mathbf{P}(\tau_x > t)$ as $t \to \infty$ or $x \to 0^+$. The function $V(x)$ is given by an explicit formula; it is the *renewal function of the ascending ladder-height process* (see Preliminaries). The case $n = 0$ has been previously studied in [17].

Theorem 1.8. (a) *If* ψ *is unbounded, and* (1.7) *and* (1.8) *hold for some* $t_0 > 0$*, then for all* $n \ge 0$ *and* $x \ge 0$*,*

$$\lim_{t \to \infty} \left(t^{n+1/2} \frac{d^n}{dt^n} \mathbf{P}(\tau_x > t) \right) = \frac{(-1)^n \Gamma(n + 1/2)}{\pi} V(x). \tag{1.12}$$

The convergence is locally uniform in $x \in [0, \infty)$*.*

(b) *If* ψ *is unbounded, regularly varying of order* $\varrho \in [0, 1]$ *at infinity, and* (1.7) *and* (1.8) *hold for some* $t_0 > 0$*, then for all* $n \ge 0$ *and* $t > t_0$ *(*$t \ge t_0$ *if* $n = 0$*),*

$$\lim_{x \to 0^+} \left(\sqrt{\psi(1/x^2)} \, \frac{d^n}{dt^n} \mathbf{P}(\tau_x > t) \right) = \frac{(-1)^n \Gamma(n + 1/2)}{\pi \Gamma(1 + \varrho)} \, \frac{1}{t^{n+1/2}} \, . \tag{1.13}$$

The convergence is uniform in $t \in (t_0, \infty)$.

Remark 1.9. The asymptotic behaviour of $(d/dt)\mathbf{P}(\tau_x > t)$ as $t \to \infty$ was completely described for (possibly asymmetric) stable Lévy processes in [14], Theorem 1. In our Theorem 1.8(a), we consider a class of symmetric, but not necessarily stable Lévy processes, and derivatives of higher order are included.

In [13], Theorem 3, an analogous problem is studied when one-dimensional distributions of the Lévy process X_t belong to the domain of attraction of a (possibly asymmetric) stable law. Again, this partially overlaps with Theorem 1.8(a), but neither result generalizes the other one. □

Finally, Theorem 1.4 is complemented by the following completeness result. It was proved in [28] under an extra assumption that the operator Π, defined in the statement of the theorem, is injective. Using methods developed partially in [29], we show that Π is always injective, and therefore the theorem holds in full generality. The present statement was announced in [28] as Theorem 1.3.

Theorem 1.10. *For* $f \in C_c((0,\infty))$ *and* $\lambda > 0$, *let*

$$\Pi f(\lambda) = \int_0^\infty f(x) F_\lambda(x) dx. \tag{1.14}$$

Then $\sqrt{2/\pi}\,\Pi$ *extends to a unitary operator on* $L^2((0,\infty))$, *which diagonalises the action of* $P_t^{(0,\infty)}$:

$$\Pi P_t^{(0,\infty)} f(\lambda) = e^{-t\psi(\lambda^2)} \Pi f(\lambda) \qquad \text{for } f \in L^2((0,\infty)).$$

Furthermore, $f \in \mathcal{D}(\mathcal{A}_{(0,\infty)}; L^2)$ *if and only if* $\psi(\lambda^2)\Pi f(\lambda)$ *is in* $L^2((0,\infty))$, *and*

$$\Pi \mathcal{A}_{(0,\infty)} f(\lambda) = -\psi(\lambda^2) \Pi f(\lambda) \qquad \text{for } f \in \mathcal{D}(\mathcal{A}_{(0,\infty)}; L^2).$$

Remark 1.11. In [28], the generalised eigenfunction expansion of Theorem 1.10 was the key step in the proof of (a restricted version of) Theorem 1.6. Here, the proofs of Theorems 1.6 and 1.10 are independent. □

We conclude the introduction with a brief description of the structure of the article. In Preliminaries, we recall the notion of complete Bernstein and Stieltjes functions, their properties and a Wiener-Hopf type transformation $\psi \mapsto \psi^\dagger$. Some new ideas are developed here, e.g. a type of continuity of the mapping $\psi \mapsto \psi^\dagger$ is proved. We also give some simple estimates related to Laplace transforms of monotone functions, and introduce the renewal function $V(x)$ of the ascending ladder-height process. In Section 3, we prove Theorem 1.10. Next two sections contain estimates and properties of ϑ_λ and $F_\lambda(x)$, respectively (see Theorem 1.4), which are essential to the derivation of the main results. In Section 6, we prove Theorems 1.6–1.8. Finally, some examples are studied in Section 7.

2. PRELIMINARIES

2.1. **Complete Bernstein functions.** A function $f(z)$ is said to be a *complete Bernstein function* (CBF in short) if

$$f(z) = c_1 + c_2 z + \frac{1}{\pi} \int_{0+}^\infty \frac{z}{z+s} \, \frac{m(ds)}{s} \,, \tag{2.1}$$

where $c_1, c_2 \ge 0$, and m is a Radon measure on $(0,\infty)$ such that $\int \min(s^{-1}, s^{-2}) m(ds) < \infty$. A function $g(z)$ is said to be a *Stieltjes function* if

$$g(z) = \frac{c_1}{z} + c_2 + \frac{1}{\pi} \int_{0+}^\infty \frac{1}{z+s} \, \tilde{m}(ds), \tag{2.2}$$

where $c_1, c_2 \geq 0$, and $\tilde{m}$ is a Radon measure on $(0, \infty)$ such that $\int \min(1, s^{-1}) \tilde{m}(ds) < \infty$.

Complete Bernstein and Stieltjes functions are often defined on $(0, \infty)$. However, (2.1) and (2.2) define holomorphic functions on $\mathbf{C} \setminus (-\infty, 0]$. We always identify functions on $(0, \infty)$ with their holomorphic extensions to $\mathbf{C} \setminus (-\infty, 0]$.

We list some basic properties of complete Bernstein and Stieltjes functions.

Proposition 2.1 (Proposition 2.18 in [28], and Corollary 6.3 in [34])**.**

(1) *Let f be a complete Bernstein function with representation* (2.1)*. Then*

$$c_1 = \lim_{z \to 0^+} f(z), \qquad c_2 = \lim_{z \to \infty} \frac{f(z)}{z}, \tag{2.3}$$

and

$$m(ds) = \lim_{\varepsilon \to 0^+} (\operatorname{Im} f(-s + i\varepsilon) ds), \tag{2.4}$$

with the limit understood in the sense of weak convergence of measures.

(2) *Let g be a Stieltjes function with representation* (2.2)*. Then*

$$c_1 = \lim_{z \to 0^+} (z g(z)), \qquad c_2 = \lim_{z \to \infty} g(z), \tag{2.5}$$

and

$$\tilde{m}(ds) + \pi c_1 \delta_0(ds) = \lim_{\varepsilon \to 0^+} (-\operatorname{Im} g(-s + i\varepsilon) ds), \tag{2.6}$$

with the limit understood in the sense of weak convergence of measures. □

Proposition 2.2 (see [34])**.** *Suppose that f and g are not constantly equal to* 0*.*

(a) *the following conditions are equivalent: $f(z)$ is CBF, $z/f(z)$ is CBF, $f(z)/z$ is Stieltjes, $1/f(z)$ is Stieltjes;*

(b) *$f(z)$ is CBF if and only if $f(z) \geq 0$ for $z \geq 0$, and f extends to a holomorphic function in $\mathbf{C} \setminus (-\infty, 0]$ such that* $\operatorname{Im} f(z) > 0$ *when* $\operatorname{Im} z > 0$*;*

(c) *if f, g are CBF and $a, c > 0$, then also $f(z) + g(z)$, $cf(z)$, $f(cz)$, $(z - a)/(f(z) - f(a))$ (extended continuously at $z = a$) and $f(g(z))$ are CBF.* □

Proposition 2.3 (Proposition 2.21 in [28])**.** *If f is a complete Bernstein function, then*

(a) *$0 \leq z f'(z) \leq f(z)$ for $z > 0$;*

(b) *$0 \leq -z f''(z) \leq 2 f'(z)$ for $z > 0$;*

(c) *$|f(z)| \leq (\sin(\varepsilon/2))^{-1} f(|z|)$ for $z \in \mathbf{C}$, $|\operatorname{Arg} z| \leq \pi - \varepsilon$, $\varepsilon \in (0, \pi)$;*

(d) *$|z f'(z)| \leq (\sin(\varepsilon/2))^{-1} f(|z|)$ for z as in (c);*

(e) *$|f(z)| \leq c(f, \varepsilon)(1 + |z|)$ for z as in (c).* □

Throughout the article, ψ usually denotes the complete Bernstein function in the Lévy-Khintchine exponent of X_t. Some preliminary results and definitions, however, are valid for more general functions ψ. If this is the case, we explicitly state all assumptions on ψ each time it is mentioned.

Following [28, 29], we define

$$\psi^\dagger(\xi) = \exp\left(\frac{1}{\pi} \int_0^\infty \frac{\xi \log \psi(\zeta^2)}{\xi^2 + \zeta^2} \, d\zeta \right) \tag{2.7}$$

for any positive function ψ for which the integral converges. In this case, $\psi^\dagger(\xi)$ is defined at least when $\operatorname{Re} \xi > 0$. By a simple substitution, for $\xi > 0$,

$$\psi^\dagger(\xi) = \exp\left(\frac{1}{\pi} \int_0^\infty \frac{\log \psi(\xi^2 \zeta^2)}{1 + \zeta^2} \, d\zeta \right). \tag{2.8}$$

Proposition 2.4 (Proposition 2.1 in [29]). *If $\psi(\xi)$ is a nonnegative function on $(0, \infty)$ and both $\psi(\xi)$ and $\xi/\psi(\xi)$ are increasing on $(0, \infty)$, then*

$$e^{-2\mathcal{C}/\pi} \sqrt{\psi(\xi^2)} \le \psi^\dagger(\xi) \le e^{2\mathcal{C}/\pi} \sqrt{\psi(\xi^2)}, \tag{2.9}$$

where $\mathcal{C} \approx 0.916$ is the Catalan constant. Note that $e^{2\mathcal{C}/\pi} \le 2$.

If, in addition, $\psi(\xi)$ is regularly varying at ∞, then

$$\lim_{\xi \to \infty} \frac{\psi^\dagger(\xi)}{\sqrt{\psi(\xi^2)}} = 1. \tag{2.10}$$

An analogous statement for $\xi \to 0$ holds for $\psi(\xi)$ regularly varying at 0.

In particular, (2.9) *holds for any CBF. Likewise,* (2.10) *holds for any regularly varying CBF.* □

The estimate (2.9) for CBFs was obtained independently in [24], Proposition 3.7, while (2.10) for CBFs was derived in [21], Proposition 2.2.

Proposition 2.5 (Proposition 3.4 in [28]). *Whenever both sides of the following identities make sense, we have:*

(a) $(1/\psi)^\dagger = 1/\psi^\dagger$, $(\psi^\varrho)^\dagger = (\psi^\dagger)^\varrho$ *($\varrho \in \mathbf{R}$) and* $(\psi_1 \psi_2)^\dagger = \psi_1^\dagger \psi_2^\dagger$*;*

(b) $(c^2 \psi)^\dagger = c \psi^\dagger$ *for $c > 0$;*

(c) *when $\psi(\xi) = \xi$, then $\psi^\dagger(\xi) = \xi$;*

(d) *if appropriate limits of ψ exist, then the corresponding limits of $\psi^\dagger$ exist, and* $\lim_{\xi \to 0^+} \psi^\dagger(\xi) = (\lim_{\xi \to 0^+} \psi(\xi))^{1/2}$, $\lim_{\xi \to \infty} \psi^\dagger(\xi) = (\lim_{\xi \to \infty} \psi(\xi))^{1/2}$. □

The first part of the following result was independently proved in [23], Proposition 2.4.

Lemma 2.6 (Lemma 3.8 in [28]). *If $\psi(\xi)$ is a CBF, then also $\psi^\dagger(\xi)$ is a CBF, and*

$$\psi^\dagger(\xi) \psi^\dagger(-\xi) = \psi(-\xi^2) \tag{2.11}$$

for $\xi \in \mathbf{C} \setminus \mathbf{R}$. □

Proposition 2.7. *Let ψ_n be a sequence of CBFs. If $\psi_n(\xi) \to \psi(\xi)$ as $n \to \infty$ for all $\xi > 0$, then $\psi_n^\dagger(\xi) \to \psi^\dagger(\xi)$ locally uniformly in $\xi \in \mathbf{C} \setminus (-\infty, 0]$.*

Proof. For any CBFs $\psi(\xi)$, $\tilde{\psi}(\xi)$, we have

$$\frac{\tilde{\psi}^\dagger(\xi)}{\psi^\dagger(\xi)} = \exp\left(\frac{1}{\pi} \int_0^\infty \frac{\xi \log(\tilde{\psi}(\zeta^2)/\psi(\zeta^2))}{\xi^2 + \zeta^2} \, d\zeta \right) \tag{2.12}$$

when $\operatorname{Re} \xi > 0$. Furthermore, by monotonicity and concavity of ψ_n, $\psi_n(1) \min(1, \zeta^2) \le \psi_n(\zeta^2) \le \psi_n(1) \max(1, \zeta^2)$ ($\zeta > 0$), and therefore there are $c_1, c_2 > 0$ such that $c_1 \min(1, \zeta^2) \le \psi_n(\zeta^2) \le c_2 \max(1, \zeta^2)$. Hence, by dominated convergence, (2.12) gives convergence of $\psi_n^\dagger(\xi)$ to $\psi^\dagger(\xi)$ when $\operatorname{Re} \xi > 0$. By Corollary 7.6(b) in [34], $\psi_n^\dagger(\xi) \to \psi^\dagger(\xi)$ for all $\xi \in \mathbf{C} \setminus (-\infty, 0]$. Pointwise convergence of CBFs is automatically locally uniform on $\mathbf{C} \setminus (-\infty, 0]$: by Proposition 2.3(d,e), when $\operatorname{Arg} \xi \in (-\pi + \varepsilon, \pi - \varepsilon)$, we have $|(\psi_n^\dagger)'(\xi)| \le c_3(\varepsilon) \psi_n^\dagger(|\xi|)/|\xi| \le c_4(\varepsilon, f) \max(|\xi|^{-1}, 1)$, which proves that $\psi_n^\dagger$ are locally equicontinuous in $\mathbf{C} \setminus (-\infty, 0]$. □

As in [28], for a nonvanishing, increasing and differentiable function ψ with strictly positive derivative, we denote

$$\psi_\lambda(\xi) = \frac{1 - \xi/\lambda^2}{1 - \psi(\xi)/\psi(\lambda^2)}, \qquad \lambda, \xi > 0, \ \lambda^2 \ne \xi. \tag{2.13}$$

This definition is extended continuously by $\psi_\lambda(\lambda^2) = \psi(\lambda^2)/(\lambda^2 \psi'(\lambda^2))$, and when ψ extends holomorphically to $\mathbf{C} \setminus (-\infty, 0]$, also $\psi_\lambda(\xi)$ is defined for $\xi \in \mathbf{C} \setminus (-\infty, 0]$. For simplicity, we denote $\psi^\dagger_\lambda(\xi) = (\psi_\lambda)^\dagger(\xi)$. By Proposition 2.2(c), if $\psi(\xi)$ is a CBF, then $\psi_\lambda(\xi)$ (and hence also $\psi^\dagger_\lambda$) is a CBF for any $\lambda > 0$.

Although (2.13) is generally used for complete Bernstein functions ψ, we will occasionaly need this definition for a *negative* function $\psi(\xi) = -\xi^\varrho$ with $\varrho \in (-1, 0)$.

Corollary 2.8. *If $\psi(\xi)$ is a CBF, then*

$$e^{-2\mathcal{C}/\pi} \sqrt{\psi_\lambda(\xi^2)} \le \psi^\dagger_\lambda(\xi) \le e^{2\mathcal{C}/\pi} \sqrt{\psi_\lambda(\xi^2)}. \qquad \square$$

Corollary 2.9. *If ψ is a CBF, then the function $\psi^\dagger_\lambda(\xi)$ is jointly continuous in $\lambda > 0$, $\xi \in \mathbf{C} \setminus (-\infty, 0)$.* □

The following monotonicity property of $\psi_\lambda(\xi)$ plays an important role.

Proposition 2.10. *Suppose that for twice differentiable, nonvanishing and increasing functions $\psi(\xi)$ and $\tilde\psi(\xi)$ we have*

$$\frac{-\psi''(\zeta)}{\psi'(\zeta)} \le \frac{-\tilde\psi''(\zeta)}{\tilde\psi'(\zeta)}, \qquad \zeta > 0. \tag{2.14}$$

Then

$$\frac{\psi_\lambda(\xi^2)}{\psi_\lambda(\lambda^2)} \le \frac{\tilde\psi_\lambda(\xi^2)}{\tilde\psi_\lambda(\lambda^2)}, \qquad 0 < \lambda < \xi,$$

$$\frac{\psi_\lambda(\xi^2)}{\psi_\lambda(\lambda^2)} \ge \frac{\tilde\psi_\lambda(\xi^2)}{\tilde\psi_\lambda(\lambda^2)}, \qquad 0 < \xi < \lambda.$$

Proof. Integration in ζ of both sides of (2.14) yields that

$$\frac{\psi'(\zeta)}{\psi'(\xi_1)} \le \frac{\tilde\psi'(\zeta)}{\tilde\psi'(\xi_1)}, \qquad 0 < \xi_1 < \zeta,$$

and by another integration in ζ,

$$\frac{\psi(\xi_2) - \psi(\xi_1)}{\psi'(\xi_1)} \le \frac{\tilde\psi(\xi_2) - \tilde\psi(\xi_1)}{\tilde\psi'(\xi_1)}, \qquad 0 < \xi_1 < \xi_2.$$

Taking $\xi_1 = \lambda^2$ and $\xi_2 = \xi^2$, we obtain

$$\frac{\psi_\lambda(\xi^2)}{\psi_\lambda(\lambda^2)} = \frac{\psi'(\lambda^2)(\lambda^2 - \xi^2)}{\psi(\lambda^2) - \psi(\xi^2)} \ge \frac{\tilde\psi'(\lambda^2)(\lambda^2 - \xi^2)}{\tilde\psi(\lambda^2) - \tilde\psi(\xi^2)} = \frac{\tilde\psi_\lambda(\xi^2)}{\tilde\psi_\lambda(\lambda^2)}, \qquad 0 < \xi < \lambda.$$

The other part is proved in a similar manner. □

Finally, recall that $\psi_\lambda(\lambda^2) = \psi(\lambda^2)/(\lambda^2 \psi'(\lambda^2))$. This gives the following simple estimate:

$$\frac{\psi_\lambda(\xi^2)}{\psi_\lambda(\lambda^2)} = \frac{\psi'(\lambda^2)(\lambda^2 - \xi^2)}{\psi(\lambda^2) - \psi(\xi^2)} \le \frac{\psi'(\lambda^2) \max(\lambda^2, \xi^2)}{|\psi(\lambda^2) - \psi(\xi^2)|}, \qquad \lambda, \xi > 0 \tag{2.15}$$

when ψ is a CBF.

Let $\psi(\xi)$ be the complete Bernstein function related with the Lévy-Khintchine exponent $\Psi(\xi)$ of the Lévy process X_t, $\Psi(\xi) = \psi(\xi^2)$. With the notation of this section, for $\lambda, \xi > 0$ we have (see Remark 4.12 in [28], and the proof of Theorem 4.1 in [29])

$$\int_0^\infty e^{-\xi x} \mathbf{P}(\tau_x > t) dx = \frac{2}{\pi} \int_0^\infty \frac{\lambda}{\lambda^2 + \xi^2} \, \frac{\lambda \psi'(\lambda^2)}{\psi(\lambda^2)} \, \psi^\dagger_\lambda(\xi) e^{-t\psi(\lambda^2)} d\lambda; \tag{2.16}$$

$$\vartheta_\lambda = \frac{1}{\pi} \int_0^\infty \frac{\lambda}{\lambda^2 - \zeta^2} \, \log \frac{\psi_\lambda(\zeta^2)}{\psi_\lambda(\lambda^2)} \, d\zeta = \operatorname{Arg} \psi^\dagger_\lambda(i\lambda); \tag{2.17}$$

$$\mathcal{L}F_\lambda(\xi) = \frac{\lambda}{\lambda^2 + \xi^2} \, \frac{\psi^\dagger_\lambda(\xi)}{\sqrt{\psi_\lambda(\lambda^2)}} \, ; \tag{2.18}$$

$$\gamma_\lambda(d\xi) = \frac{1}{\pi} \frac{\sqrt{\psi_\lambda(\lambda^2)}}{\psi^\dagger_\lambda(\xi)} \, \operatorname{Im} \frac{\lambda \psi'(\lambda^2)}{\psi(\lambda^2) - \psi^+(-\xi^2)} \, d\xi; \tag{2.19}$$

for the last equality the assumption of the final part of Theorem 1.4 is required.

2.2. **Estimates for the Laplace transform.** This short section contains some rather standard estimates for the inverse Laplace transform, similar to those used in [29].

Proposition 2.11. *Let $a > 0$, $c \ge 1$. If f is nonnegative and $f(x) \le c f(a) \max(1, x/a)$ ($x > 0$), then for any $\xi > 0$,*

$$f(a) \ge \frac{\xi \mathcal{L}f(\xi)}{c(1 + (a\xi)^{-1} e^{-a\xi})}.$$

Proof. We have

$$\begin{aligned} \xi \mathcal{L}f(\xi) &= \int_0^a \xi e^{-\xi x} f(x) dx + \int_a^\infty \xi e^{-\xi x} f(x) dx \\ &\le c f(a) \int_0^a \xi e^{-\xi x} dx + \frac{c f(a)}{a} \int_a^\infty \xi x e^{-\xi x} dx \\ &= c f(a)(1 - e^{-a\xi}) + \frac{c f(a)}{a\xi} (1 + a\xi) e^{-a\xi} \\ &= c f(a)(1 + (a\xi)^{-1} e^{-a\xi}). \end{aligned}$$

□

Proposition 2.12. *If f is nonnegative and increasing, then for $a, \xi > 0$,*

$$f(a) \le e^{a\xi} \xi \mathcal{L}f(\xi).$$

Proof. As before,

$$\begin{aligned} \xi \mathcal{L}f(\xi) &= \int_0^a \xi e^{-\xi x} f(x) dx + \int_a^\infty \xi e^{-\xi x} f(x) dx \\ &\ge f(a) \int_a^\infty \xi e^{-\xi x} dx = f(a) e^{-a\xi}. \end{aligned}$$

□

Proposition 2.13. *Let $b \ge a > 0$, and suppose that f is nonnegative and increasing on $(0, b)$, and $f(x) \ge m$ for $x \ge b$. Then for any $\xi > 0$,*

$$f(a) \le \frac{e^{a\xi} \xi \mathcal{L}f(\xi) - m e^{-(b-a)\xi}}{1 - e^{-(b-a)\xi}}.$$

Proof. By Proposition 2.12 applied to $f(x) \mathbf{1}_{(0,a)}(x) + f(a) \mathbf{1}_{[a,\infty)}(x)$,

$$\begin{aligned} f(a) &\le e^{a\xi} \xi \left(\int_0^\infty e^{-\xi x} f(x) dx + \int_a^\infty e^{-\xi x} (f(a) - f(x)) dx \right) \\ &\le e^{a\xi} \xi \left(\int_0^\infty e^{-\xi x} f(x) dx + \int_b^\infty e^{-\xi x} (f(a) - m) dx \right) \\ &= e^{a\xi} \xi \mathcal{L}f(\xi) + e^{-(b-a)\xi} (f(a) - m). \end{aligned}$$

□

2.3. **Elements of fluctuation theory.** In the sequel we will need the function $V(x)$, which is described by its Laplace transform, $\mathcal{L}V(\xi) = \xi/\psi^\dagger(\xi)$. By Proposition 2.2(a) and Lemma 2.6, $\xi/\psi^\dagger(\xi)$ is a complete Bernstein function. Hence, V is a Bernstein function, i.e. V is nonnegative and V' is completely monotone (see [34]). By Theorem 4.4 in [29],

$$\frac{1}{5}\,\frac{1}{\sqrt{\psi(1/x^2)}} \le V(x) \le 5\,\frac{1}{\sqrt{\psi(1/x^2)}}\,, \qquad x > 0.$$

Suppose that ψ extends to a function ψ^+ holomorphic in the upper complex half-plane and continuous in the region $\operatorname{Im}\xi \ge 0$. Then by Proposition 4.5 in [29],

$$V(x) = bx + \frac{1}{\pi}\int_{0^+}^\infty \operatorname{Im}\left(-\frac{1}{\psi^+(-\xi^2)}\right)\frac{\psi^\dagger(\xi)}{\xi}\,(1-e^{-x\xi})d\xi, \qquad x > 0,$$

where $b = (\lim_{\xi\to 0^+}(\xi/\psi(\xi)))^{1/2}$.

Probabilistically, $V(x)$ is the renewal function for the ascending ladder-height process H_s corresponding to X_t. When ψ is unbounded, then X_t satisfies the absolute continuity condition and $V(x)$ is the (unique up to a multiplicative constant) increasing harmonic function for X_t on $(0,\infty)$, cf. [35].

3. Eigenfunction expansion

In this section we prove a peculiar integral identity, which (together with the results of [28]) yields a short proof of Theorem 1.10.

Lemma 3.1. *Let $\psi(\xi)$ be an arbitrary positive, continuously differentiable function on $(0,\infty)$ with strictly positive derivative, and such that $(\log(1+\psi(\zeta^2)))/\zeta^2$ is integrable on $(0,\infty)$. Then for all $\xi_1, \xi_2 > 0$,*

$$\int_0^\infty \frac{\lambda^4\psi'(\lambda^2)(\xi_1+\xi_2)\psi_\lambda^\dagger(\xi_1)\psi_\lambda^\dagger(\xi_2)}{\psi(\lambda^2)(\lambda^2+\xi_1^2)(\lambda^2+\xi_2^2)}\,d\lambda = \frac{\pi}{2}. \tag{3.1}$$

Proof. As in the proof of Theorem 4.1 in [29], for $\xi > 0$ and $z \in \mathbf{C}\setminus(-\infty,0]$ we define

$$\varphi(\xi,z) = \exp\left(-\frac{1}{\pi}\int_0^\infty \frac{\xi\log(1+\psi(\zeta^2)/z)}{\xi^2+\zeta^2}\,d\zeta\right),$$

where log is the principal branch of the logarithm. Clearly, for each $\xi > 0$, $\varphi(\xi,z)$ is a holomorphic function of z, and $\varphi(\xi,z) > 0$ when $z > 0$. As it was observed in [29], when $\operatorname{Im} z > 0$, we have $\operatorname{Arg}(1+\psi(\zeta^2)/z) \in (-\pi,0)$, so that

$$\operatorname{Arg}\varphi(\xi,z) = -\frac{1}{\pi}\int_0^\infty \frac{\xi\operatorname{Arg}(1+\psi(\zeta^2)/z)}{\xi^2+\zeta^2}\,d\zeta \in (0,\pi/2).$$

Hence, for any $\xi_1,\xi_2 > 0$, $\operatorname{Im}(\varphi(\xi_1,z)\varphi(\xi_2,z)) > 0$ when $\operatorname{Im} z > 0$. By Proposition 2.2(b), $f(z) = \varphi(\xi_1,z)\varphi(\xi_2,z)$ is a CBF of z.

By monotone convergence, $f(z)$ converges to 0 when $z \to 0^+$, and by dominated convergence, $\lim_{z\to\infty} f(z) = 1$. It follows that the constants c_1, c_2 in the representation (2.1) for the CBF $f(z)$ vanish. Furthermore, for any $\xi > 0$, $\varphi(\xi,z)$ extends to a continuous function $\varphi^+(\xi,z)$ in the region $\operatorname{Im} z \ge 0$, given by the formula

$$\varphi^+(\xi,z) = \exp\left(-\frac{1}{\pi}\int_0^\infty \frac{\xi\log^-(1+\psi(\zeta^2)/z)}{\xi^2+\zeta^2}\,d\zeta\right),$$

where $\log^-(z)$ is the continuous extension of $\log z$ to the region $\operatorname{Im} z \le 0$. The measure m in the representation (2.1) for the function $f(z) = \varphi(\xi_1,z)\varphi(\xi_2,z)$ is therefore given by

$m(ds) = \pi^{-1} \operatorname{Im}(\varphi^+(\xi_1, -s)\varphi^+(\xi_2, -s))ds$ (Proposition 2.1(a)). By (2.1) and monotone convergence,

$$\begin{aligned} &\frac{1}{\pi} \int_0^\infty \frac{\operatorname{Im}(\varphi^+(\xi_1, -s)\varphi^+(\xi_2, -s))}{s} \, ds \\ &\qquad = \frac{1}{\pi} \lim_{z \to \infty} \int_0^\infty \frac{z \operatorname{Im}(\varphi^+(\xi_1, -s)\varphi^+(\xi_2, -s))}{z + s} \, \frac{ds}{s} \\ &\qquad = \lim_{z \to \infty} (\varphi(\xi_1, z)\varphi(\xi_2, z)) = 1. \end{aligned} \tag{3.2}$$

Note that if $\psi(\xi)$ is bounded on $(0, \infty)$ and $s \ge \sup_{\xi > 0} \psi(\xi)$, then $\varphi^+(\xi, -s)$ is real, so that the integrand in (3.2) vanishes for such s. Hence, we may substitute $\psi(\lambda^2)$ for s in (3.2) to obtain that

$$\frac{2}{\pi} \int_0^\infty \lambda \psi'(\lambda^2) \operatorname{Im} \frac{\varphi^+(\xi_1, -\psi(\lambda^2))\varphi^+(\xi_2, -\psi(\lambda^2))}{\psi(\lambda^2)} \, d\lambda = 1.$$

Finally, in the proof of Theorem 4.1 in [29] (formula (4.3) therein) it is proved that

$$\varphi^+(\xi, -\psi(\lambda^2)) = \frac{\lambda(\lambda + \xi i)\psi_\lambda^\dagger(\xi)}{\lambda^2 + \xi^2} \, .$$

Since $\operatorname{Im}((\lambda + \xi_1 i)(\lambda + \xi_2 i)) = \lambda(\xi_1 + \xi_2)$, the lemma is proved. □

Remark 3.2. Lemma 3.1 was conjectured in a preliminary version of [28]. □

Proof of Theorem 1.10. In [28], Theorem 1.10 was proved under an extra assumption that the operator Π defined in (1.14) is injective. Below we show that this condition is always satisfied.

Let $e_\xi(x) = e^{-\xi x} \mathbf{1}_{(0,\infty)}(x)$ ($\xi > 0$). By (2.18), for $\lambda > 0$,

$$\Pi e_\xi(\lambda) = \int_0^\infty e^{-\xi x} F_\lambda(x) dx = \mathcal{L} F_\lambda(\xi) = \sqrt{\frac{\lambda^2 \psi'(\lambda^2)}{\psi(\lambda^2)}} \, \frac{\lambda}{\lambda^2 + \xi^2} \, \psi_\lambda^\dagger(\xi).$$

By Lemma 3.1,

$$\begin{aligned} \langle \Pi e_{\xi_1}, \Pi e_{\xi_2} \rangle_{L^2((0,\infty))} &= \int_0^\infty \frac{\lambda^4 \psi'(\lambda^2) \psi_\lambda^\dagger(\xi_1) \psi_\lambda^\dagger(\xi_2)}{\psi(\lambda^2)(\lambda^2 + \xi_1^2)(\lambda^2 + \xi_2^2)} \, d\lambda \\ &= \frac{\pi}{2(\xi_1 + \xi_2)} = \frac{\pi}{2} \, \langle e_{\xi_1}, e_{\xi_2} \rangle_{L^2((0,\infty))} \, . \end{aligned}$$

The functions e_ξ ($\xi > 0$) form a linearly dense set in $L^2((0, \infty))$. By approximation, $\sqrt{2/\pi} \, \Pi$ is a unitary operator on $L^2((0, \infty))$. In particular, Π is injective. □

4. Estimates of ϑ_λ

As it will become clear in the next section, upper bounds for the phase shift ϑ_λ (which is defined by (1.5)) are crucial for applications of Theorems 1.4 and 1.10: when ϑ_λ is close to $\pi/2$, estimates of the eigenfunctions $F_\lambda(x)$ are problematic. Recall that by (1.5) and (2.17),

$$\vartheta_\lambda = \operatorname{Arg} \psi_\lambda^\dagger(i\lambda) = -\frac{1}{\pi} \int_0^\infty \frac{1}{1 - \zeta^2} \log \frac{\psi_\lambda(\lambda^2 \zeta^2)}{\psi_\lambda(\lambda^2)} \, d\zeta, \qquad \lambda > 0. \tag{4.1}$$

Here $\psi(\xi)$ is a complete Bernstein function in the Lévy-Khintchine exponent of X_t. By a substitution $\zeta = 1/s$ in the integral over $(1, \infty)$, one obtains that (see Proposition 4.16

in [28])

$$\vartheta_\lambda = \frac{1}{\pi} \int_0^1 \frac{1}{1 - \zeta^2} \log \frac{\psi_\lambda(\lambda^2/\zeta^2)}{\psi_\lambda(\lambda^2\zeta^2)} \, d\zeta, \qquad \lambda > 0. \tag{4.2}$$

When $\psi(\xi) = \xi^{\alpha/2}$ ($0 < \alpha \le 2$), that is, when X_t is the symmetric α-stable process, we have $\vartheta_\lambda = (2 - \alpha)\pi/8$ (Example 6.1 in [28]). In the general case, $0 \le \vartheta_\lambda < \pi/2$, and $\vartheta_\lambda \le (\pi/2) \sup_{\xi>0}(\xi\psi'_\lambda(\xi)/\psi_\lambda(\xi))$ (Proposition 4.18 in [28]). This estimate is insufficient for our needs, and it is significantly improved below. By (4.1) and Proposition 2.10, we have the following monotonicity result. Note that ψ need not be a complete Bernstein function in Propositions 4.1 and 4.2.

Proposition 4.1. *Suppose that ψ and $\tilde{\psi}$ are arbitrary twice-differentiable, increasing and nonvanishing functions (not necessarily complete Bernstein functions). Define ϑ_λ by (4.2), and define $\tilde{\vartheta}_\lambda$ in a similar way, using $\tilde{\psi}$. We assume that the integrals in the definitions of ϑ_λ and $\tilde{\vartheta}_\lambda$ are convergent and finite. If $-\psi''(\xi)/\psi'(\xi) \le -\tilde{\psi}''(\xi)/\tilde{\psi}'(\xi)$ for all $\xi > 0$, then $\vartheta_\lambda \le \tilde{\vartheta}_\lambda$ for all $\lambda > 0$.* ☐

Taking $\tilde{\psi}(\xi) = \xi^{\alpha/2}$ (corresponding to the symmetric α-stable process) and using the fact that $\tilde{\vartheta}_\lambda = (2 - \alpha)\pi/8$, we obtain a global estimate for ϑ_λ. It turns out that although this method works for $\alpha \in (0, 2]$, it generalises easily to $\alpha \in [-2, 2]$. First, we recall the following result, which was given in Example 6.1 in [28] for $\varrho \in (0, 1]$; the proof, however, works for any $\varrho \ne 0$.

Proposition 4.2 (see Example 6.1 in [28])**.** *Let $\varrho \ne 0$ and $\psi(\xi) = \xi^\varrho$ if $\varrho > 0$, $\psi(\xi) = -\xi^\varrho$ if $\varrho < 0$. Define ϑ_λ by (4.2). Then $\vartheta_\lambda = (1 - \varrho)\pi/4$.* ☐

From now on, ψ is again the complete Bernstein function such that $\Psi(\xi) = \psi(\xi^2)$ is the Lévy-Khintchine exponent of the process X_t.

Proposition 4.3. *We have*

$$\left(\inf_{\xi>0} \frac{\xi|\psi''(\xi)|}{\psi'(\xi)} \right) \frac{\pi}{4} \le \vartheta_\lambda \le \left(\sup_{\xi>0} \frac{\xi|\psi''(\xi)|}{\psi'(\xi)} \right) \frac{\pi}{4}. \tag{4.3}$$

Proof. Denote the supremum in (4.3) by $1 - \varrho$. By Proposition 2.3(b), $\varrho \in [-1, 1]$. Suppose that ϱ is non-zero, and let $\tilde{\psi}(\xi) = \xi^\varrho$ if $\varrho > 0$, $\tilde{\psi}(\xi) = -\xi^\varrho$ if $\varrho < 0$. Observe that $\tilde{\psi}$ is increasing and $-\xi\tilde{\psi}''(\xi)/\tilde{\psi}'(\xi) = (1 - \varrho)$, so that

$$\frac{-\psi''(\zeta)}{\psi'(\zeta)} \le \frac{-\tilde{\psi}''(\zeta)}{\tilde{\psi}'(\zeta)}, \qquad \zeta > 0.$$

The upper bound in (4.3) follows by Propositions 4.1 and 4.2. When $\varrho = 0$, we simply consider $\tilde{\psi}(\xi) = -\xi^{-\varepsilon}$ and let $\varepsilon \to 0^+$. Finally, the lower bound is proved in a similar manner. ☐

We conclude this section with a *local* estimate of ϑ_λ, which depends solely on $\psi(\lambda^2)$, $\psi'(\lambda^2)$ and $\psi''(\lambda^2)$. This result is used only in the construction of an irregular example in Subsection 7.6. We need the following technical result.

Proposition 4.4. *For $a > 0$,*

$$-\frac{1}{\pi} \int_0^\infty \frac{1}{1 - x^2} \log \frac{1 + a^2x^2}{1 + a^2} \, dx = \arctan a.$$

For $b \in (0, 1)$,

$$-\frac{1}{\pi} \int_0^1 \frac{1}{1 - x^2} \log(1 - b^2(1 - x^2)) dx = \frac{(\arcsin b)^2}{\pi}.$$

Proof. The function $\log((1+a^2x^2)/(1+a^2))$ is holomorphic in the upper half-plane with a branch cut along $(ia^{-1}, i\infty)$. By an appropriate contour integration and limit procedure (we omit the details),

$$\begin{aligned} 2\int_0^\infty \frac{1}{1-x^2} \log \frac{1+a^2x^2}{1+a^2}\, dx &= \int_{-\infty}^\infty \frac{1}{1-x^2} \log \frac{1+a^2x^2}{1+a^2}\, dx \\ &= \int_{ia^{-1}}^{i\infty} \frac{2\pi i}{1-x^2}\, dx = -2\pi \int_{1/a}^\infty \frac{1}{1+y^2}\, dy = -2\pi \arctan a. \end{aligned}$$

For the second equality, we have by Taylor expansion and beta integral,

$$\begin{aligned} -\int_0^\infty \frac{1}{1-x^2} \log(1-b^2(1-x^2)) dx &= \sum_{n=1}^\infty \frac{b^{2n}}{n} \int_0^1 (1-x^2)^{n-1} dx \\ &= \sum_{n=1}^\infty \frac{b^{2n}}{2n} \int_0^1 (1-s)^{n-1} s^{-1/2} ds = \sum_{n=1}^\infty \frac{\Gamma(n)\Gamma(1/2) b^{2n}}{2n\Gamma(n+1/2)} = (\arcsin b)^2; \end{aligned}$$

see e.g. p. 108 in [31] for the last identity. □

Proposition 4.5. *We have*

$$\vartheta_\lambda \le \frac{\pi}{2} - \arcsin \sqrt{\frac{\lambda^2 \psi'(\lambda^2)}{\psi(\lambda^2)}}$$

and

$$\begin{aligned} \vartheta_\lambda \ge \left(\arcsin \sqrt{\frac{\lambda^2 |\psi''(\lambda^2)|}{2\psi'(\lambda^2)}} \right)^2 + \left(\arcsin \sqrt{\frac{\lambda^2 \psi(\lambda^2) |\psi''(\lambda^2)|}{2\psi'(\lambda^2)(\psi(\lambda^2) - \lambda^2\psi'(\lambda^2))}} \right)^2 \\ - \left(\arcsin \sqrt{\frac{\lambda^4 |\psi''(\lambda^2)|}{2(\psi(\lambda^2) - \lambda^2\psi'(\lambda^2))}} \right)^2 . \end{aligned}$$

It is easy to see that these bounds are sharp for $\psi(\xi) = c_1\xi/(c_2+\xi)$ and $\psi(\xi) = c\xi$ (and, in fact, these are all CBFs ψ for which equalities hold), see examples in Section 7.

Proof. Note that $\psi_\lambda(0) = 1$ and $\psi_\lambda(\lambda^2) = \psi(\lambda^2)/(\lambda^2\psi'(\lambda^2))$. Let $a^2 = \psi_\lambda(\lambda^2) - 1$. Since ψ_λ is concave, we have $\psi_\lambda(\lambda^2\zeta^2) \ge 1 + a^2\zeta^2$ for $\zeta < 1$, and $\psi_\lambda(\lambda^2\zeta^2) \le 1 + a^2\zeta^2$ for $\zeta > 1$. Hence, by Proposition 4.4,

$$\begin{aligned} \vartheta_\lambda &= -\frac{1}{\pi} \int_0^\infty \frac{1}{1-\zeta^2} \log \frac{\psi_\lambda(\lambda^2\zeta^2)}{\psi_\lambda(\lambda^2)}\, d\zeta \\ &\le -\frac{1}{\pi} \int_0^\infty \frac{1}{1-\zeta^2} \log \frac{1+a^2\zeta^2}{1+a^2}\, d\zeta = \arctan a. \end{aligned}$$

Since $\arctan a = \pi/2 - \arcsin(1/(1+a^2)^{1/2})$, this proves the upper bound.

For the lower bound, let $a_1^2 = \psi_\lambda(\lambda^2) = 1 + a^2$ and $a_2^2 = \lambda^2\psi_\lambda'(\lambda^2)/\psi_\lambda(\lambda^2)$. By a short calculation, $a_2^2 = \lambda^2|\psi''(\lambda^2)|/(2\psi'(\lambda^2))$. Since $\psi_\lambda(\xi)$ is concave, its graph lies below the tangent line at $\xi = \lambda^2$, so that

$$\begin{aligned} \psi_\lambda(\lambda^2\zeta^2) &\le \psi_\lambda(\lambda^2) - \lambda^2\psi_\lambda'(\lambda^2)(1-\zeta^2) \\ &= a_1^2(1 - a_2^2(1-\zeta^2)). \end{aligned}$$

A similar argument for the CBF $\xi/(\psi_\lambda(\xi)-1)$ and a short calculation give

$$\begin{aligned}\frac{\xi}{\psi_\lambda(\xi)-1} &\le \frac{\lambda^2}{a^2} + \frac{a^2-a_1^2a_2^2}{a^4}\,(\xi-\lambda^2)\\ &= \frac{a^2\xi - a_1^2a_2^2(\xi-\lambda^2)}{a^4}\,,\end{aligned}$$

which for $\xi = \lambda^2/\zeta^2$ ($\zeta > 0$) reduces to

$$\begin{aligned}\psi_\lambda(\lambda^2/\zeta^2) &\ge 1 + \frac{a^4\zeta^{-2}}{a^2\zeta^{-2} - a_1^2a_2^2(\zeta^{-2}-1)}\\ &= \frac{a_1^2(a^2-a_2^2(1-\zeta^2))}{a^2+a_1^2a_2^2(1-\zeta^2)}\,.\end{aligned}$$

Therefore, by (4.2),

$$\begin{aligned}\vartheta_\lambda &= \frac{1}{\pi}\int_0^1 \frac{1}{1-\zeta^2}\log\frac{\psi_\lambda(\lambda^2/\zeta^2)}{\psi_\lambda(\lambda^2\zeta^2)}\,d\zeta\\ &\ge \frac{1}{\pi}\int_0^1 \frac{1}{1-\zeta^2}\log\frac{1-a^{-2}a_2^2(1-\zeta^2)}{(1-a_2^2(1-\zeta^2))(1-a^{-2}a_1^2a_2^2(1-\zeta^2))}\,d\zeta.\end{aligned}$$

Triple application of Proposition 4.4 yields

$$\pi\vartheta_\lambda \ge (\arcsin a_2)^2 + (\arcsin(a_1a_2/a))^2 - (\arcsin(a_2/a))^2. \qquad \square$$

5. PROPERTIES OF $F_\lambda(x)$

Some basic estimates of the eigenfunctions $F_\lambda(x)$ have already been established in [28]. However, for the proof of Theorems 1.6–1.8, more detailed properties of $F_\lambda(x)$ are required. Recall that by (1.3) and (2.18),

$$\mathcal{L}F_\lambda(\xi) = \frac{\lambda}{\lambda^2+\xi^2}\,\frac{\psi_\lambda^\dagger(\xi)}{\sqrt{\psi_\lambda(\lambda^2)}}, \qquad \lambda > 0,\ \operatorname{Re}\xi > 0, \tag{5.1}$$

and by (1.4),

$$F_\lambda(x) = \sin(\lambda x + \vartheta_\lambda) - G_\lambda(x), \qquad \lambda > 0,\ x > 0.$$

The phase shift ϑ_λ was studied in detail in the previous section. Recall that the correction term $G_\lambda(x)$ is a completely monotone function, $G_\lambda(x) = \mathcal{L}\gamma_\lambda(x)$, and γ_λ is a finite measure on $(0,\infty)$. In many important cases, γ_λ is given explicitly by (1.6) and (2.19); however, in this section these identities are not used. We extend the definition of $F_\lambda(x)$ by letting $F_\lambda(x) = 0$ for $x \le 0$.

It was proved in Proposition 4.22 in [28] that

$$\mathcal{L}F_\lambda(\xi) \le \frac{\lambda+\xi}{\lambda^2+\xi^2} \le \frac{2}{\lambda+\xi}, \qquad \lambda,\xi > 0. \tag{5.2}$$

By combining (5.1) and Proposition 2.4, we obtain the following more detailed estimate.

Corollary 5.1. *We have*

$$e^{-2\mathcal{C}/\pi}\,\frac{\lambda}{\lambda^2+\xi^2}\sqrt{\frac{\psi_\lambda(\xi^2)}{\psi_\lambda(\lambda^2)}} \le \mathcal{L}F_\lambda(\xi) \le e^{2\mathcal{C}/\pi}\,\frac{\lambda}{\lambda^2+\xi^2}\sqrt{\frac{\psi_\lambda(\xi^2)}{\psi_\lambda(\lambda^2)}} \tag{5.3}$$

for $\lambda,\xi > 0$. $\square$

Hölder continuity of F_λ was already studied in Lemma 4.24 in [28]. However, when ψ does not have a power-type growth at infinity, F_λ fails to be Hölder continuous. On the other hand, uniform estimate for $F_\lambda(x)$ for small $x > 0$ is crucial in the proof of Theorem 1.6. We generalize the estimates of [28] in the two lemmas below.

Since $G_\lambda(x)$ is completely monotone, we have $G'_\lambda(x) \le 0$ and $G''_\lambda(x) \ge 0$ for $\lambda, x > 0$. It follows that the function $F_\lambda(x) = \sin(\lambda x + \vartheta_\lambda) - G_\lambda(x)$ is increasing on $[0, (\pi/2 - \vartheta_\lambda)/\lambda]$ and concave on $[0, (\pi - \vartheta_\lambda)/\lambda]$ (this explains the role of upper bounds for ϑ_λ). Furthermore, $0 \le G_\lambda(x) \le \sin\vartheta_\lambda$ (Lemma 4.21 in [28]), and so $-1 - \sin\vartheta_\lambda \le F_\lambda(x) \le 1$.

Lemma 5.2. *When $\lambda > 0$ and $0 < \lambda x \le (\pi/2 - \vartheta_\lambda)/2$, we have*

$$\begin{aligned} \frac{e^{1-2\mathcal{C}/\pi}}{2(e+1)} \lambda x \sqrt{\frac{\psi_\lambda(1/x^2)}{\psi_\lambda(\lambda^2)}} &\le F_\lambda(x) \\ &\le \left(2e^{1+2\mathcal{C}/\pi} + \frac{4e}{\pi/2 - \vartheta_\lambda}\right) \lambda x \sqrt{\frac{\psi_\lambda(1/x^2)}{\psi_\lambda(\lambda^2)}}. \end{aligned} \tag{5.4}$$

Proof. Let $\lambda > 0$, $b = (\pi/2 - \vartheta_\lambda)/\lambda$ and $0 < a \le b$. By concavity of the sine function and convexity of G_λ, for any $x > a$ we have

$$\frac{\sin(\lambda x + \vartheta_\lambda) - \sin\vartheta_\lambda}{x} \le \frac{\sin(\lambda a + \vartheta_\lambda) - \sin\vartheta_\lambda}{a},$$
$$\frac{\sin\vartheta_\lambda - G_\lambda(x)}{x} \le \frac{\sin\vartheta_\lambda - G_\lambda(a)}{a}.$$

Hence $F_\lambda(x) \le (x/a) F_\lambda(a)$ when $x > a$. This and monotonicity of F_λ on $[0, b]$ yield

$$F_\lambda(x) \le F_\lambda(a) \max(1, x/a)$$

for all $x > 0$. By Proposition 2.11, for $\xi > 0$ we have

$$F_\lambda(a) \ge \frac{\xi \mathcal{L}F_\lambda(\xi)}{1 + (a\xi)^{-1} e^{-a\xi}}.$$

For $\xi = 1/a$, this gives

$$F_\lambda(a) \ge \frac{e}{e+1} \, \frac{\mathcal{L}F_\lambda(1/a)}{a}.$$

On the other hand, $F_\lambda(x)$ is increasing in $x \in (0, b)$, and clearly $F_\lambda(x) \ge -2$ for all $x \ge b$. Proposition 2.13 gives

$$F_\lambda(a) \le \frac{e^{a\xi} \xi \mathcal{L}F_\lambda(\xi) + 2e^{-(\pi/2 - \vartheta_\lambda - \lambda a)\xi/\lambda}}{1 - e^{-(\pi/2 - \vartheta_\lambda - \lambda a)\xi/\lambda}}$$

for all $\xi > 0$. Let $\xi = 1/a$. Using the inequality $1 - e^{-s} \ge s/(s+1)$ $(s > 0)$ in the denominator and $e^{-1/s} \le s$ $(s > 0)$ in the numerator, we obtain that

$$\begin{aligned} F_\lambda(a) &\le \frac{1}{1 - e^{-(\pi/2 - \vartheta_\lambda - \lambda a)/(\lambda a)}} \left(\frac{e\mathcal{L}F_\lambda(1/a)}{a} + 2e^{-(\pi/2 - \vartheta_\lambda - \lambda a)/(\lambda a)} \right) \\ &= \frac{e}{1 - e^{-(\pi/2 - \vartheta_\lambda - \lambda a)/(\lambda a)}} \left(\frac{\mathcal{L}F_\lambda(1/a)}{a} + 2e^{-(\pi/2 - \vartheta_\lambda)/(\lambda a)} \right) \\ &\le \frac{e(\pi/2 - \vartheta_\lambda)}{\pi/2 - \vartheta_\lambda - \lambda a} \left(\frac{\mathcal{L}F_\lambda(1/a)}{a} + \frac{2\lambda a}{\pi/2 - \vartheta_\lambda} \right). \end{aligned}$$

The above bounds for $F_\lambda(a)$ combined with (5.3) give

$$\begin{aligned} \frac{e^{1-2\mathcal{C}/\pi}}{e+1} \, \frac{\lambda/a}{\lambda^2+1/a^2} \sqrt{\frac{\psi_\lambda(1/a^2)}{\psi_\lambda(\lambda^2)}} &\le F_\lambda(a) \\ &\le \frac{e(\pi/2-\vartheta_\lambda)}{\pi/2-\vartheta_\lambda-\lambda a} \left(e^{2\mathcal{C}/\pi} \, \frac{\lambda/a}{\lambda^2+1/a^2} \sqrt{\frac{\psi_\lambda(1/a^2)}{\psi_\lambda(\lambda^2)}} + \frac{2\lambda a}{\pi/2-\vartheta_\lambda} \right) . \end{aligned}$$

Assume now that $a \in (0, b/2]$. Then $a \le b/2 = (\pi/2 - \vartheta_\lambda)/(2\lambda) \le 1/\lambda$, and therefore $1/a^2 \le \lambda^2 + 1/a^2 \le 2/a^2$. It follows that

$$\begin{aligned} \frac{e^{1-2\mathcal{C}/\pi}\lambda a}{2(e+1)} \sqrt{\frac{\psi_\lambda(1/a^2)}{\psi_\lambda(\lambda^2)}} &\le F_\lambda(a) \\ &\le \frac{e(\pi/2-\vartheta_\lambda)}{\pi/2-\vartheta_\lambda-\lambda a} \left(e^{2\mathcal{C}/\pi} \lambda a \sqrt{\frac{\psi_\lambda(1/a^2)}{\psi_\lambda(\lambda^2)}} + \frac{2\lambda a}{\pi/2-\vartheta_\lambda} \right) . \end{aligned}$$

Since $\psi_\lambda(\xi)$ is increasing on $(0, \infty)$ and $\lambda < 1/a$, we have $1 \le \sqrt{\psi_\lambda(1/a^2)/\psi_\lambda(\lambda^2)}$. Finally, $\lambda a \le \lambda b/2 = (\pi/2 - \vartheta_\lambda)/2$, so that $(\pi/2 - \vartheta_\lambda)/(\pi/2 - \vartheta_\lambda - \lambda a) \le 2$. Formula (5.4) follows. □

Since $F_\lambda(x) = \sin(\lambda x + \vartheta_\lambda) - G_\lambda(x)$ for a completely monotone G_λ, the modulus of continuity of F_λ is described by the behavior of $F_\lambda(x)$ for small x. More precisely, we have the following result.

Lemma 5.3. *We have*

$$|F_\lambda(x) - F_\lambda(y)| \le \min\left(\frac{30\lambda|x-y|}{\pi/2-\vartheta_\lambda} \sqrt{\frac{\psi_\lambda(1/|x-y|^2)}{\psi_\lambda(\lambda^2)}} \, , 4 \right) \tag{5.5}$$

for all $\lambda > 0$ *and* $x, y \ge 0$.

Proof. The inequality $|F_\lambda(x) - F_\lambda(y)| \le 4$ is clear. For $x, y > 0$ and $\lambda > 0$,

$$\begin{aligned} |F_\lambda(x) - F_\lambda(y)| &\le |G_\lambda(x) - G_\lambda(y)| + \lambda|x-y| \\ &\le |G_\lambda(|x-y|) - G_\lambda(0^+)| + \lambda|x-y| \\ &\le |F_\lambda(|x-y|)| + 2\lambda|x-y|. \end{aligned} \tag{5.6}$$

The same inequality is obviously true also when $x = 0$ or $y = 0$, so from now on we assume that $x, y \ge 0$.

Suppose that $\lambda|x-y| \le (\pi/2 - \vartheta_\lambda)/2$. Then, by (5.6) and (5.4),

$$|F_\lambda(x) - F_\lambda(y)| \le \left(2e^{1+2\mathcal{C}/\pi} + \frac{4e}{\pi/2-\vartheta_\lambda} \right) \lambda|x-y| \sqrt{\frac{\psi_\lambda(1/|x-y|^2)}{\psi_\lambda(\lambda^2)}} + 2\lambda|x-y|.$$

As in the proof of Lemma 5.2, $1 \le \sqrt{\psi_\lambda(1/|x-y|^2)/\psi_\lambda(\lambda^2)}$ (indeed $\lambda < 1/|x-y|$, and ψ_λ is increasing), so that

$$|F_\lambda(x) - F_\lambda(y)| \le \left(2e^{1+2\mathcal{C}/\pi} + \frac{4e}{\pi/2-\vartheta_\lambda} + 2 \right) \lambda|x-y| \sqrt{\frac{\psi_\lambda(1/|x-y|^2)}{\psi_\lambda(\lambda^2)}}.$$

Finally, the parenthesised expression does not exceed $(\pi e^{1+2\mathcal{C}/\pi} + 4e + \pi)/(\pi/2 - \vartheta_\lambda)$, and (5.5) follows.

It remains to consider $\lambda|x-y| > (\pi/2 - \vartheta_\lambda)/2$. We claim that in this case, the minimum in (5.5) is equal to 4. By Proposition 2.2(a), $f(\xi) = \xi/\psi_\lambda(\xi)$ is a CBF. Hence, $f(x)/f(a) \ge \min(1, x/a)$. We obtain

$$\begin{aligned}\frac{\lambda|x-y|}{\pi/2-\vartheta_\lambda}\sqrt{\frac{\psi_\lambda(1/|x-y|^2)}{\psi_\lambda(\lambda^2)}} &= \frac{1}{\pi/2-\vartheta_\lambda}\sqrt{\frac{\lambda^2/\psi_\lambda(\lambda^2)}{(1/|x-y|^2)/\psi_\lambda(1/|x-y|^2)}} \\ &\ge \frac{\min(\lambda|x-y|,1)}{\pi/2-\vartheta_\lambda} > \frac{1}{2}.\end{aligned}$$

This proves our claim, and the proof is complete. □

The behavior of $F_\lambda(x)$ when $x \to 0^+$ (with fixed $\lambda > 0$) was studied in Lemma 4.27 in [28]: if ψ is an unbounded CBF regularly varying of order $\alpha \in [0,1]$ at infinity, then $F_\lambda(x)$ is regularly varying of order α at 0, and

$$\lim_{x\to0^+}\Big(\sqrt{\psi(1/x^2)}\,F_\lambda(x)\Big) = \frac{\sqrt{\lambda^2\psi'(\lambda^2)}}{\Gamma(1+\alpha)}, \qquad \lambda > 0. \tag{5.7}$$

The next result shows that as $\lambda \to 0^+$ with fixed $x > 0$, $F_\lambda(x)$ behaves as $\lambda\sqrt{\psi'(\lambda^2)}V(x)$, where $V(x)$ is the renewal function of the ascending ladder-height process, $\mathcal{L}V(\xi) = \xi/\psi^\dagger(\xi)$ (see Preliminaries).

Proposition 5.4. *Suppose that ψ is an unbounded CBF. The function $F_\lambda(x)$ is jointly continuous in $\lambda > 0$, $x \ge 0$. Furthermore, if $\limsup_{\lambda\to0^+}\vartheta_\lambda < \pi/2$, then*

$$\lim_{\lambda\to0^+}\frac{F_\lambda(x)}{\lambda\sqrt{\psi'(\lambda^2)}} = V(x), \qquad x \ge 0,$$

and the convergence is locally uniform in $x \ge 0$. In other words, $F_\lambda(x)/(\lambda\sqrt{\psi'(\lambda^2)})$ extends to a continuous function in $\lambda, x \ge 0$.

Proof. Roughly speaking, we use estimates of $\mathcal{L}F_\lambda$ to show L^2 continuity of $e^{-x}F_\lambda(x)$ in $\lambda > 0$, and then estimates of F_λ to replace L^2 convergence by locally uniform convergence.

Let $\lambda_n \to \lambda > 0$. By Corollary 2.9, $\mathcal{L}F_{\lambda_n}(1+i\xi)$ converges to $\mathcal{L}F_\lambda(1+i\xi)$ pointwise for $\xi \in \mathbf{R}$. By (5.2) and dominated convergence, $\mathcal{L}F_{\lambda_n}(1+i\xi)$ converges to $\mathcal{L}F_\lambda(1+i\xi)$ in $L^2(\mathbf{R})$. Note that $\mathcal{L}F_\lambda(1+i\xi)$ is the Fourier transform of $e^{-x}F_\lambda(x)$. By Plancherel's theorem, $e^{-x}F_{\lambda_n}(x)$ converges to $e^{-x}F_\lambda(x)$ in $L^2(\mathbf{R})$. By (5.5), the sequence $F_{\lambda_n}(x)$ is equicontinuous in x (here we use the assumption that ψ is unbounded), and hence it converges to $F_\lambda(x)$ locally uniformly in $x \ge 0$. The first part of the proposition is proved.

As $\lambda \to 0^+$, the functions $(\lambda^2/\psi(\lambda^2))\psi_\lambda(\xi)$ converge pointwise to the CBF $\xi/\psi(\xi)$. Therefore, by Proposition 2.7, $\sqrt{\lambda^2/\psi(\lambda^2)}\,\psi_\lambda^\dagger(\xi)$ converges to $\xi/\psi^\dagger(\xi)$ ($\xi \in \mathbf{C}\setminus(-\infty,0]$). We conclude that $\mathcal{L}F_\lambda(1+i\xi)/(\lambda\sqrt{\psi'(\lambda^2)})$ converges to $\mathcal{L}V(1+i\xi)$ ($\xi \in \mathbf{R}$). Again, we obtain $L^2(\mathbf{R})$ convergence of $e^{-x}F_\lambda(x)/(\lambda\sqrt{\psi'(\lambda^2)})$ to $e^{-x}V(x)$. The proof will be complete if we show that the family of functions $F_\lambda(x)/(\lambda\sqrt{\psi'(\lambda^2)})$ is equicontinuous as $\lambda \to 0^+$.

By (5.5), we have

$$\begin{aligned}\frac{|F_\lambda(x)-F_\lambda(y)|}{\lambda\sqrt{\psi'(\lambda^2)}} &\le \frac{30|x-y|}{\pi/2-\vartheta_\lambda}\sqrt{\frac{\psi_\lambda(1/|x-y|^2)}{\psi'(\lambda^2)\psi_\lambda(\lambda^2)}} \\ &= \frac{30}{\pi/2-\vartheta_\lambda}\sqrt{\frac{1-\lambda^2|x-y|^2}{\psi(1/|x-y|^2)-\psi(\lambda^2)}}.\end{aligned} \tag{5.8}$$

Fix $\lambda_0 > 0$ small enough. By the assumption, $\pi/2 - \vartheta_\lambda$ is bounded below by a positive constant for $\lambda \in (0, \lambda_0)$. It follows that for $\lambda \in (0, \lambda_0)$ and $x, y \ge 0$ satisfying $|x - y| < 1/\lambda_0$, the expression on the right hand side of (5.8) is bounded above by $c/\sqrt{\psi(1/|x-y|)^2 - \psi(\lambda_0^2)}$ (with c depending only on ψ and λ_0). This upper bound does not depend on $\lambda \in (0, \lambda_0)$, and since ψ is unbounded, it converges to 0 as $|x-y| \to 0$. □

6. SUPREMA AND FIRST PASSAGE TIMES FOR COMPLETE BERNSTEIN FUNCTIONS

Below we prove Theorems 1.6–1.8. We remark that the condition (1.7) is used only to assert that $\sup_{\lambda>0} \vartheta_\lambda < \pi/2$, and can be replaced by the latter condition.

Proof of Theorem 1.6. First we consider $n = 0$, that is, we will show that

$$\mathbf{P}(\tau_x > t) = \frac{2}{\pi} \int_0^\infty \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t\psi(\lambda^2)} F_\lambda(x) d\lambda. \tag{6.1}$$

We claim that the integral in (6.1) converges, and that $e^{-\xi x} e^{-t\psi(\lambda^2)} F_\lambda(x) \sqrt{\psi'(\lambda^2)/\psi(\lambda^2)}$ is jointly integrable in $\lambda, x > 0$ for any $\xi > 0$. Assuming this is true, the proof is quite straightforward. Indeed, by Fubini, the Laplace transform (in x) of the right hand side of (6.1) is then

$$\frac{2}{\pi} \int_0^\infty \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t\psi(\lambda^2)} \mathcal{L}F_\lambda(\xi) d\lambda.$$

By Theorem 1.3 (see (2.16)) and Theorem 1.4 (see (2.18)), this is equal to the Laplace transform (in x) of $\mathbf{P}(\tau_x > t)$, and the result follows by the uniqueness of the Laplace transform. Hence it is enough to prove our claim.

Let $t \ge t_0$, $\Theta = \sup_{\lambda>0} \vartheta_\lambda$ and $\lambda_0 = (\pi/2 - \Theta)/(2x)$. By the assumption (1.7) and Proposition 4.3, $\Theta < \pi/2$, and hence $\lambda_0 > 0$. Note that $\lambda_0 x \le \pi/4 < 1$. Since $|F_\lambda(x)| \le 2$ and $\sqrt{\psi'(\lambda^2)/\psi(\lambda^2)} \le 1/\lambda$, we have

$$\begin{aligned}
&\int_{\lambda_0}^\infty \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t\psi(\lambda^2)} |F_\lambda(x)| d\lambda \\
&\qquad \le 2 \int_{\min(\lambda_0, 1)}^1 \frac{e^{-t\psi(\lambda^2)}}{\lambda}\, d\lambda + 2 \int_1^\infty \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t\psi(\lambda^2)} d\lambda \\
&\qquad \le 2 \max(0, -\log \lambda_0) + 2 \int_1^\infty \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t_0\psi(\lambda^2)} d\lambda,
\end{aligned}$$

which is finite by assumption (1.8). We now consider $\lambda \in (0, \lambda_0)$. By Lemma 5.2, we have

$$0 \le \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t\psi(\lambda^2)} F_\lambda(x) \le c_1(\Theta) \lambda x \sqrt{\frac{\psi'(\lambda^2)\psi_\lambda(1/x^2)}{\psi(\lambda^2)\psi_\lambda(\lambda^2)}}\, .$$

Furthermore, by (2.15) (recall that $\lambda < \lambda_0 < 1/x$),

$$\lambda x \sqrt{\frac{\psi'(\lambda^2)\psi_\lambda(1/x^2)}{\psi(\lambda^2)\psi_\lambda(\lambda^2)}} \le \frac{\lambda \psi'(\lambda^2)}{\sqrt{\psi(\lambda^2)(\psi(1/x^2) - \psi(\lambda^2))}}\, .$$

Integration and substitution $z = \psi(\lambda^2)$ give

$$\begin{aligned}\int_0^{\lambda_0} \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t\psi(\lambda^2)} |F_\lambda(x)| d\lambda &\le c_1(\Theta) \int_0^{\lambda_0} \frac{\lambda \psi'(\lambda^2)}{\sqrt{\psi(\lambda^2)(\psi(1/x^2) - \psi(\lambda^2))}}\, d\lambda \\ &= \frac{c_1(\Theta)}{2} \int_0^{\psi(\lambda_0^2)} \frac{1}{\sqrt{z(\psi(1/x^2) - z)}}\, dz \le \frac{c_1(\Theta)\pi}{2}\,.\end{aligned}$$

We conclude that

$$\int_0^\infty \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t\psi(\lambda^2)} |F_\lambda(x)| d\lambda \le c_2(\psi, t) + 2\max(0, \log x),$$

which shows that the integral in (6.1) is absolutely convergent. This also shows that

$$\int_0^\infty \int_0^\infty e^{-\xi x} \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}}\, e^{-t\psi(\lambda^2)} |F_\lambda(x)| d\lambda d\xi < \infty,$$

as desired. The proof in the case $n = 0$ is complete.

For general $n \ge 0$, one uses dominated convergence to prove that the derivative can by taken under the integral sign; we omit the details. □

We now find upper and lower bounds for the derivatives in t of $\mathbf{P}(\tau_x > t)$. Note that the estimates of $\mathbf{P}(\tau_x > t)$ are covered by [29] for a wider class of Lévy processes. We begin with a simple technical result.

Proposition 6.1. (a) *Let $\gamma(a; x) = \int_0^x e^{-s} s^{a-1} ds$ $(a, x > 0)$ be the lower incomplete gamma function. Then*

$$\frac{\min(1, x^a)}{ae} \le \gamma(a; x) \le \frac{\min(\Gamma(a+1), x^a)}{a}\,, \qquad a, x > 0. \tag{6.2}$$

(b) *We have*

$$\int_0^x \frac{e^{-s} s^{a-1}}{\sqrt{x - s}}\, ds \le \frac{\min(2\Gamma(a) + \pi a^a e^{-a}, 2(1/a + 1)x^a)}{\sqrt{2x}}\,, \qquad a, x > 0. \tag{6.3}$$

Proof. For the part (a), we simply have

$$\gamma(a; x) = \int_0^x e^{-s} s^{a-1} ds \ge \frac{1}{e} \int_0^{\min(1,x)} s^{a-1} ds = \frac{\min(1, x^a)}{ae}\,,$$

and

$$\gamma(a; x) = \int_0^x e^{-s} s^{a-1} ds \le \min\left(\Gamma(a), \int_0^x s^{a-1} ds\right) = \min\left(\Gamma(a), \frac{x^a}{a}\right).$$

To prove (b), we split the integral into two parts. First,

$$\begin{aligned}\int_0^{x/2} \frac{e^{-s} s^{a-1}}{\sqrt{x - s}}\, ds &\le \sqrt{\frac{2}{x}} \int_0^{x/2} e^{-s} s^{a-1} ds \\ &= \sqrt{\frac{2}{x}}\, \gamma(a; x/2) \le \sqrt{\frac{2}{x}}\, \min\left(\Gamma(a), \frac{x^a}{a}\right).\end{aligned}$$

Next, $s^a e^{-s}$ attains its maximum at $s = a$. Hence,

$$\begin{aligned}\int_{x/2}^{x} \frac{e^{-s} s^{a-1}}{\sqrt{x-s}} \, ds &\le a^a e^{-a} \int_{x/2}^{x} \frac{1}{\sqrt{s^2(x-s)}} \, ds \\ &\le a^a e^{-a} \sqrt{\frac{2}{x}} \int_{x/2}^{x} \frac{1}{\sqrt{s(x-s)}} \, ds = \frac{\pi a^a e^{-a}}{2} \sqrt{\frac{2}{x}} \, .\end{aligned}$$

Finally,

$$\int_{x/2}^{x} \frac{e^{-s} s^{a-1}}{\sqrt{x-s}} \, ds \le x^{a-1} \int_{x/2}^{x} \frac{1}{\sqrt{x-s}} \, ds = \sqrt{2} \, x^{a-1/2}.$$

It follows that

$$\begin{aligned}\int_{0}^{x} \frac{e^{-s} s^{a-1}}{\sqrt{x-s}} \, ds &\le \sqrt{\frac{2}{x}} \, \frac{\min(\Gamma(a+1), x^a)}{a} + \min\left(\frac{\pi a^a e^{-a}}{2} \sqrt{\frac{2}{x}}, \sqrt{2} \, x^{a-1/2} \right) \\ &\le \sqrt{\frac{2}{x}} \left(\frac{\min(\Gamma(a+1), x^a)}{a} + \min\left(\frac{\pi a^a e^{-a}}{2}, x^a \right) \right) ,\end{aligned}$$

which gives (6.3). ☐

Lemma 6.2. *Suppose that* (1.7) *and* (1.8) *hold for some* $t_0 > 0$. *For* $n \ge 0$, $t > t_0$ *(*$t \ge t_0$ *if* $n = 0$*) and* $\lambda_0 > 0$, *denote (cf* (1.8)*)*

$$I_n(t_0, \lambda_0) = \frac{4}{\pi} \int_{\lambda_0}^{\infty} e^{-t_0 \psi(\lambda^2)} \sqrt{\psi'(\lambda^2)} (\psi(\lambda^2))^{n-1/2} d\lambda.$$

Let $\Theta = \sup_{\xi > 0} \vartheta_\lambda$ *and* $\lambda_0(x) = (\pi/2 - \Theta)/(2x)$. *If* $t > t_0$ *(*$t \ge t_0$ *if* $n = 0$*) and* $x > 0$, *then*

$$\begin{aligned} & C_1(n, \Theta) \min \left((\psi(1/x^2))^n, \frac{1}{t^{n+1/2} \sqrt{\psi(1/x^2)}} \right) + I_n(t, \lambda_0(x)) \\ & \qquad \le (-1)^n \, \frac{d^n}{dt^n} \, \mathbf{P}(\tau_x > t) \qquad (6.4) \\ & \qquad \le C_2(n, \Theta) \min \left((\psi(1/x^2))^n, \frac{1}{t^{n+1/2} \sqrt{\psi(1/x^2)}} \right) - I_n(t, \lambda_0(x)). \end{aligned}$$

Remark 6.3.

1. When (1.8) holds for some $t_0 > 0$, then clearly $I_0(t_0, \lambda_0) < \infty$ for any $\lambda_0 > 0$. Furthermore, for any $n > 0$, $t > t_0$ and $\lambda_0 > 0$, we have $e^{-(t-t_0)\psi(\lambda^2)} \le c(n, \lambda_0)(\psi(\lambda^2))^n$ for $\lambda \ge \lambda_0$. Hence, (1.8) implies also that $I_n(t, \lambda_0) < \infty$ for any $n > 0$, $t > t_0$ and $\lambda_0 > 0$.
2. When $t > t_0$, then $I_n(t, \lambda_0) \le e^{-(t-t_0)\psi(\lambda_0^2)} I_n(t_0, \lambda_0)$. Hence, $I_n(t, \lambda_0)$, if finite, decays exponentially fast as $t \to \infty$. Therefore, $I_n(t, \lambda_0)$ can be considered as the 'error term'; see Theorem 1.8. ☐

Proof of Lemma 6.2. By the assumption (1.7) and Proposition 4.3, $\Theta < \pi/2$. Let $k = (\pi/2 - \Theta)/2 \in (0, \pi/4)$, so that $\lambda_0 = \lambda_0(x) = k/x$. Denote the integrand in (1.9) by $f_{n,t,x}(\lambda)$,

$$f_{n,t,x}(\lambda) = \sqrt{\frac{\psi'(\lambda^2)}{\psi(\lambda^2)}} \, (\psi(\lambda^2))^n e^{-t\psi(\lambda^2)} F_\lambda(x).$$

By Lemma 5.2, for $\lambda \in (0, \lambda_0)$ we have

$$\begin{aligned} f_{n,t,x}(\lambda) &\ge (\psi(\lambda^2))^{n-1/2} \sqrt{\psi'(\lambda^2)}\, e^{-t\psi(\lambda^2)}\, \frac{e^{1-2\mathcal{C}/\pi} \lambda x}{2(e+1)} \sqrt{\frac{\psi_\lambda(1/x^2)}{\psi_\lambda(\lambda^2)}} \\ &= \frac{e^{1-2\mathcal{C}/\pi} \lambda \psi'(\lambda^2)}{2(e+1)}\, e^{-t\psi(\lambda^2)} \frac{(\psi(\lambda^2))^{n-1/2} \sqrt{1-\lambda^2 x^2}}{\sqrt{\psi(1/x^2) - \psi(\lambda^2)}} \\ &\ge \frac{e^{1-2\mathcal{C}/\pi} \lambda \psi'(\lambda^2)}{2(e+1)}\, e^{-t\psi(\lambda^2)} \frac{(\psi(\lambda^2))^{n-1/2} \sqrt{1-k^2}}{\sqrt{\psi(1/x^2))}} \,. \end{aligned}$$

Since $\lambda x < k$ and $\psi(\lambda_0^2) = \psi(k^2/x^2) \ge k^2 \psi(1/x^2)$ (ψ is nonnegative and concave, and $k < 1$), we obtain (with $z = \psi(\lambda^2)$)

$$\begin{aligned} \int_0^{\lambda_0} f_{n,t,x}(\lambda) d\lambda &\ge \frac{e^{1-2\mathcal{C}/\pi} \sqrt{1-k^2}}{4(e+1)\sqrt{\psi(1/x^2)}} \int_0^{\psi(\lambda_0^2)} z^{n-1/2} e^{-tz} dz \\ &\ge \frac{e^{1-2\mathcal{C}/\pi} \sqrt{1-k^2}}{4(e+1) t^{n+1/2} \sqrt{\psi(1/x^2)}}\, \gamma(n+1/2; k^2 t \psi(1/x^2)), \end{aligned}$$

where γ is the lower incomplete gamma function. By (6.2),

$$\int_0^{\lambda_0} f_{n,t,x}(\lambda) d\lambda \ge \frac{e^{1-2\mathcal{C}/\pi} \sqrt{1-k^2}}{4(e+1) t^{n+1/2} \sqrt{\psi(1/x^2)}}\, \frac{\min(1, (k^2 t \psi(1/x^2))^{n+1/2})}{(n+1/2)e} \,. \tag{6.5}$$

In a similar way,

$$f_{n,t,x}(\lambda) \le \left(2e^{1+2\mathcal{C}/\pi} + \frac{2e}{k}\right) \lambda \psi'(\lambda^2) e^{-t\psi(\lambda^2)}\, \frac{(\psi(\lambda^2))^{n-1/2}}{\sqrt{\psi(1/x^2) - \psi(\lambda^2)}},$$

and so

$$\begin{aligned} \int_0^{\lambda_0} f_{n,t,x}(\lambda) d\lambda &\le \left(e^{1+2\mathcal{C}/\pi} + \frac{e}{k}\right) \int_0^{\psi(1/x^2)} \frac{z^{n-1/2} e^{-tz}}{\sqrt{\psi(1/x^2) - z}}\, dz \\ &= \left(e^{1+2\mathcal{C}/\pi} + \frac{e}{k}\right) \frac{1}{t^n} \int_0^{t\psi(1/x^2)} \frac{s^{n-1/2} e^{-s}}{\sqrt{t\psi(1/x^2) - s}}\, ds. \end{aligned}$$

By (6.3), with $c_1(n) = 2\Gamma(n+1/2) + \pi((n+1/2)/e)^{n+1/2}$ and $c_2(n) = 2(1/(n+1/2)+1)$,

$$\int_0^{\lambda_0} f_{n,t,x}(\lambda) d\lambda \le \left(e^{1+2\mathcal{C}/\pi} + \frac{e}{k}\right) \frac{\min(c_1(n), c_2(n)(t\psi(1/x^2))^{n+1/2})}{t^n \sqrt{2t\psi(1/x^2)}}. \tag{6.6}$$

Hence, we found a two-sided estimate for the integral of $f_{n,t,x}(\lambda)$ over $(0, \lambda_0)$. The integral over (λ_0, ∞) is highly oscillatory, and therefore difficult to estimate. For this reason, we are satisfied with a simple bound obtained using the inequality $|F_\lambda(x)| \le 2$,

$$\begin{aligned} \frac{2}{\pi} \int_{\lambda_0}^\infty |f_{n,t,x}(\lambda)| d\lambda &\le \frac{4}{\pi} \int_{\lambda_0}^\infty (\psi(\lambda^2))^{n-1/2} \sqrt{\psi'(\lambda^2)} e^{-t\psi(\lambda^2)} d\lambda \\ &= I_n(t, \lambda_0). \end{aligned} \tag{6.7}$$

The lower bound in (6.4) is a consequence of (6.5) and (6.7), and the upper bound in (6.4) follows from (6.6) and (6.7). □

We remark that in the statement of the lemma, we can take

$$C_1(n,\Theta) = \frac{e^{1-2\mathcal{C}/\pi}\sqrt{1-\pi^2/16}\,(\pi/4-\Theta/2)^{2n+1}}{2\pi e(e+1)(n+1/2)} \ge \frac{(\pi/2-\Theta)^{2n+1}}{17(n+1/2)2^{2n+3}}\,,$$

$$C_2(n,\Theta) = \frac{e\sqrt{2}}{\pi}\left(e^{2\mathcal{C}/\pi} + \frac{2}{\pi/2-\Theta}\right)\bigl(2\Gamma(n+1/2) + \pi((n+1/2)/e)^{n+1/2}\bigr) \le \frac{15\,n!}{\pi/2-\Theta}\,.$$

Note that $C_1(n,\Theta)$ decreases with Θ, while $C_2(n,\Theta)$ increases with Θ. The notation of Lemma 6.2, namely $I_n(t_0,\lambda_0)$, $C_1(n,\Theta)$ and $C_2(n,\Theta)$, is kept in the remaining part of the section.

Corollary 6.4. *Let* $\varepsilon > 0$, $n \ge 1$, $t_0 > 0$, $x_0 > 0$. *If the conditions* (1.7) *and* (1.8) *hold true, then there are positive constants* $C_3(n,\Theta)$, $C_4(n,\Theta)$, $C_5(\varepsilon,n,\Theta,I_n(t_0,\lambda_0(x_0)))$ *(here* $\Theta = \sup_{\xi>0}\vartheta_\lambda$ *and* $\lambda_0(x) = (\pi/2-\Theta)/(2x)$*) such that*

$$\frac{C_3(n,\Theta)}{t^{n+1/2}\sqrt{\psi(1/x^2)}} \le (-1)^n\,\frac{d^n}{dt^n}\,\mathbf{P}(\tau_x > t) \le \frac{C_4(n,\Theta)}{t^{n+1/2}\sqrt{\psi(1/x^2)}} \tag{6.8}$$

when $x \in (0,x_0]$ *and*

$$t \ge \max\left((1+\varepsilon)t_0, \frac{1}{\psi(1/x^2)}, \frac{C_5(\varepsilon,n,\Theta,I_n(t_0,\lambda_0(x_0)))}{(\psi(1/x^2))^{1+\varepsilon}}\right).$$

Proof. This is a combination of Lemma 6.2 and Remark 6.3. As in the proof of Lemma 6.2, we let $k = (\pi/2-\Theta)/2$, so that $\lambda_0 = \lambda_0(x) = k/x$. Recall that since $k < 1$, we have $\psi(k^2/x^2) \ge k^2\psi(1/x^2)$. It follows that for $n \ge 1$, $\varepsilon > 0$, $x > 0$ and $t > (1+\varepsilon)t_0$, we have

$$\begin{aligned}\frac{I_n(t,\lambda_0)}{I_n(t_0,\lambda_0)} &\le e^{-(t-t_0)\psi(\lambda_0^2)} \le e^{-\varepsilon/(1+\varepsilon)t\psi(k^2/x^2)} \\ &\le e^{-k^2\varepsilon/(1+\varepsilon)t\psi(1/x^2)} \le \frac{c_2(\varepsilon,n,\Theta)}{(t\psi(1/x^2))^{n+1/2+n/\varepsilon}}\,.\end{aligned}$$

Fix $A > 0$. When $t(\psi(1/x^2))^{1+\varepsilon} \ge A^\varepsilon$, we obtain

$$I_n(t,\lambda_0) \le \frac{c_2(\varepsilon,n,\Theta)}{A^n t^{n+1/2}\sqrt{\psi(1/x^2)}}\,I_n(t_0,\lambda_0).$$

Hence, by (6.4), if $t\psi(1/x^2) \ge 1$, we have (with the constants $C_1(n,\Theta)$ and $C_2(n,\Theta)$ of Lemma 6.2)

$$\begin{aligned}\left(C_1(n,\Theta) - \frac{c_2(\varepsilon,n,\Theta)I_n(t_0,\lambda_0)}{A^n}\right)\frac{1}{t^{n+1/2}\sqrt{\psi(1/x^2)}} &\le (-1)^n\,\frac{d^n}{dt^n}\,\mathbf{P}(\tau_x > t) \\ &\le \left(C_2(n,\Theta) + \frac{c_2(\varepsilon,n,\Theta)I_n(t_0,\lambda_0)}{A^n}\right)\frac{1}{t^{n+1/2}\sqrt{\psi(1/x^2)}}.\end{aligned}$$

When $x \in (0,x_0]$, then $I_n(t_0,\lambda_0(x)) \le I_n(t_0,\lambda_0(x_0))$. Hence, (6.8) holds if $A > (C_5(\varepsilon,n,\Theta,I_n(t_0,\lambda_0(x_0))))^{1/\varepsilon}$ for some C_5. ☐

Proof of Theorem 1.7. Part (a) follows directly from Corollary 6.4. For part (b), suppose that $|\psi''(\xi)|/\psi'(\xi) \le \varrho/\xi$ for some $\varrho \in [0,1)$ and all $\xi > 0$. By Proposition 4.3, with the notation of Lemma 6.2, we have $\Theta \le \varrho\pi/4$.

Integrating the inequality $-\psi''(\xi)/\psi'(\xi) \le \varrho/\xi$, we obtain $\psi'(\xi_1)/\psi'(\xi_2) \le (\xi_2/\xi_1)^\varrho$ if $0 < \xi_1 < \xi_2$. Integrating this again in ξ_1 gives $\psi(\xi)/\psi'(\xi) \le \xi/(1-\varrho)$. Hence, for all

$t > 0$ and $\lambda_0 > 0$,

$$\begin{aligned} I_n(t, \lambda_0) &\le \frac{4}{\pi} \int_{\lambda_0}^{\infty} e^{-t\psi(\lambda^2)} \, \frac{\lambda \psi'(\lambda^2)}{\sqrt{(1 - \varrho)\psi(\lambda^2)}} \, (\psi(\lambda^2))^{n-1/2} d\lambda \\ &= \frac{2}{\pi \sqrt{1 - \varrho} \, t^n} \int_{\lambda_0}^{\infty} e^{-t\psi(\lambda^2)} (t\psi(\lambda^2))^{n-1} (2\lambda t \psi'(\lambda^2)) d\lambda \\ &= \frac{2}{\pi \sqrt{1 - \varrho} \, t^n} \, \Gamma(n; t\psi(\lambda_0^2)), \end{aligned}$$

where $\Gamma(a; z)$ is the upper incomplete Gamma function. In particular, $I_n(t, \lambda_0)$ is finite, and (1.8) holds true.

Let $k = (\pi/2 - \Theta)/2 \in (\pi/8, \pi/4)$, and take $\lambda_0 = \lambda_0(x) = k/x$, as in Lemma 6.2. Recall that since $k < 1$, we have $\psi(\lambda_0^2) \ge k^2 \psi(1/x^2)$. Hence,

$$\begin{aligned} I_n(t, \lambda_0) &\le \frac{2}{\pi \sqrt{1 - \varrho}} \, \frac{\Gamma(n; k^2 t \psi(1/x^2))}{t^n} \\ &= \frac{2}{\pi \sqrt{1 - \varrho}} \, \frac{\sqrt{t\psi(1/x^2)} \, \Gamma(n; (\pi/8)^2 t \psi(1/x^2))}{t^{n+1/2} \sqrt{\psi(1/x^2)}} \, . \end{aligned} \tag{6.9}$$

The constant $c_3(n, \varrho)$ in the statement of the theorem is so chosen that for all $s \ge c_3(n, \varrho)$,

$$\frac{2}{\pi \sqrt{1 - \varrho}} \, \sqrt{s} \, \Gamma(n; (\pi/8)^2 s) \le \frac{\min(C_1(n, \pi/4), C_2(n, \pi/4))}{2} \, ,$$

with the constants $C_1(n, \Theta)$ and $C_2(n, \Theta)$ from Lemma 6.2. Then, by Lemma 6.2, when $t\psi(1/x^2) \ge c_3(n, \varrho)$ we have

$$\frac{C_1(n, \pi/4)}{2 t^{n+1/2} \sqrt{\psi(1/x^2)}} \le (-1)^n \, \frac{d^n}{dt^n} \, \mathbf{P}(\tau_x > t) \le \frac{C_2(n, \pi/4)}{2 t^{n+1/2} \sqrt{\psi(1/x^2)}},$$

as desired. Finally, by (6.9), we have $I_n(t, \lambda_0) \le c(n, \varrho)/(t^{n+1/2} \sqrt{\psi(1/x^2)})$, where $c(n, \varrho) = 2\pi^{-1} (1 - \varrho)^{-1/2} \sup_{s > 0} (s^{1/2} \Gamma(n; (\pi/8)^2 s))$. This and Lemma 6.2 prove that the upper bound in (1.11) holds for all $t, x > 0$, but with a constant depending on ϱ. □

Remark 6.5. (a) The strict inequality $\varrho < 1$ is essential to the proof. The case $\varrho = 1$ is much more complicated; see the example in Subsection 7.3.

(b) Theorem 1.7(b) can be easily generalised to the case when $|\psi''(\xi)|/\psi'(\xi) \le \varrho/\xi$ only for $\xi \in (0, \varepsilon) \cup (1/\varepsilon, \infty)$, with $\varepsilon \in (0, 1)$ fixed. In this case the constant $c_3 = c_3(n, \varrho, \varepsilon)$ depends also on ε. We omit the details.

(c) The upper bound in (1.11) is certainly not optimal when $t\psi(1/x^2)$ is small. This is due to essential cancellations in (1.9). □

Proof of Theorem 1.8(a). We use the notation of Lemma 6.2, and take $\lambda_0 = \lambda_0(x) = (\pi/2 - \Theta)/(2x)$ for a fixed $x > 0$. Let $f_{n,t,x}(\lambda)$ be the integrand in (1.9). We have

$$t^{n+1/2} \int_0^{\lambda_0} f_{n,t,x}(\lambda) d\lambda = \int_0^{\lambda_0} \frac{F_\lambda(x)}{\lambda \sqrt{\psi'(\lambda^2)}} \, t^{n+1/2} (\psi(\lambda^2))^{n-1/2} e^{-t\psi(\lambda^2)} \lambda \psi'(\lambda^2) d\lambda.$$

By Proposition 5.4, $\lim_{\lambda \to 0} F_\lambda(x)/(\lambda \sqrt{\psi'(\lambda^2)}) = V(x)$, and therefore $F_\lambda(x)/(\lambda \sqrt{\psi'(\lambda^2)})$ extends to a continuous function of $\lambda \in [0, \lambda_0]$. Let

$$\mu_t(d\lambda) = t^{n+1/2} (\psi(\lambda^2))^{n-1/2} e^{-t\psi(\lambda^2)} \lambda \psi'(\lambda^2) \mathbf{1}_{[0,\lambda_0]}(\lambda) d\lambda.$$

As $t \to \infty$, the density function of μ_t converges uniformly to 0 on $[\varepsilon, \lambda_0]$ for every $\varepsilon > 0$. Hence, $\mu_t(d\lambda)$ converges weakly to a point-mass at 0. Furthermore, by a substitution

$z = t\psi(\lambda^2)$,

$$\|\mu_t\| = \frac{1}{2} \int_0^{t\psi(\lambda_0^2)} z^{n-1/2} e^{-z} dz = \frac{\gamma(n+1/2; t\psi(\lambda_0^2))}{2},$$

and hence $\|\mu_t\|$ converges to $\Gamma(n+1/2)/2$ as $t \to \infty$. It follows that

$$\lim_{t\to\infty} \left(t^{n+1/2} \int_0^{\lambda_0} f_{n,t,x}(\lambda) d\lambda \right) = \frac{\Gamma(n+1/2)}{2} V(x).$$

Finally, by (6.7) and Remark 6.3,

$$\lim_{t\to\infty} \left| t^{n+1/2} \frac{4}{\pi} \int_{\lambda_0}^{\infty} f_{n,t,x}(\lambda) d\lambda \right| \le \lim_{t\to\infty} \left(t^{n+1/2} I_n(t, \lambda_0) \right) = 0,$$

and so (1.12) follows by (1.9). The convergence is locally uniform, since the extension of $F_\lambda(x)/(\lambda\sqrt{\psi'(\lambda^2)})$ is jointly continuous in $\lambda \ge 0$ and $x \ge 0$ (Proposition 5.4), and $t^{n+1/2} I_n(t, \lambda_0(x))$ converges to 0 locally uniformly in $x \ge 0$. ☐

Remark 6.6. Alternatively, one can deduce formula (1.12) (without uniformity in x) as follows. In [17] it was proved that $\sqrt{t}\,\mathbf{P}(\tau_x > t)$ converges to $V(x)/\sqrt{\pi}$ as $t \to \infty$. By Theorem 1.7(a), $\mathbf{P}(\tau_x > t)$ is ultimately completely monotone. As it was observed in [14], Remark 4, this already implies formula (1.12) for all $n \ge 0$, see [8] (we omit the details). ☐

Proof of Theorem 1.8(b). The argument is similar to the proof of part (a) of the theorem. Again we use the notation of the proof of Lemma 6.2. Let $f_{n,t,x}(\lambda)$ be the integrand in (1.9). Fix $t > t_0$. We have

$$\begin{aligned} &\sqrt{\psi(1/x^2)} \int_0^\infty f_{n,t,x}(\lambda) d\lambda \\ &\qquad = \int_0^\infty \frac{\sqrt{\psi(1/x^2)}\, F_\lambda(x)}{\lambda\sqrt{\psi'(\lambda^2)}} (\psi(\lambda^2))^{n-1/2} e^{-t\psi(\lambda^2)} \lambda \psi'(\lambda^2) d\lambda. \end{aligned}$$

By (5.7), $\lim_{x\to 0^+} \sqrt{\psi(1/x^2)}\, F_\lambda(x)/(\lambda\sqrt{\psi'(\lambda^2)}) = 1/\Gamma(1+\alpha)$. We will use dominated convergence for the integral over an initial interval $(0, B)$, and a simple uniform bound on the remaining interval $[B, \infty)$.

Let $B > 0$. Consider x small enough, so that $\lambda x \le (\pi/2 - \Theta)/2$ and $\psi(1/x^2) \ge 2\psi(\lambda^2)$ for $\lambda \in (0, B)$ (recall that ψ is unbounded). By Lemma 5.2, for $\lambda \in (0, B)$,

$$\begin{aligned} \frac{\sqrt{\psi(1/x^2)}\, F_\lambda(x)}{\lambda\sqrt{\psi'(\lambda^2)}} &\le c(\Theta) \sqrt{\frac{\lambda^2 x^2 \psi(1/x^2) \psi_\lambda(1/x^2)}{\psi'(\lambda^2)\psi_\lambda(\lambda^2)}} \\ &= c(\Theta) \sqrt{\frac{\psi(1/x^2)(1 - \lambda^2 x^2)}{\psi(1/x^2) - \psi(\lambda^2)}} \\ &\le c(\Theta) \frac{1}{\sqrt{1 - \psi(\lambda^2)/\psi(1/x^2)}} \le \sqrt{2}\, c(\Theta). \end{aligned}$$

Hence, by dominated convergence,

$$\begin{aligned} &\lim_{x\to 0^+} \left(\sqrt{\psi(1/x^2)} \int_0^B f_{n,t,x}(\lambda) d\lambda \right) \\ &\qquad = \frac{1}{\Gamma(1+\alpha)} \int_0^B (\psi(\lambda^2))^{n-1/2} e^{-t\psi(\lambda^2)} \lambda \psi'(\lambda^2) d\lambda. \end{aligned}$$

More precisely, we have

$$\lim_{x\to 0^+}\int_0^B \left|\sqrt{\psi(1/x^2)}f_{n,t,x}(\lambda) - \frac{(\psi(\lambda^2))^{n-1/2}e^{-t\psi(\lambda^2)}\lambda\psi'(\lambda^2)}{\Gamma(1+\alpha)}\right| d\lambda = 0,$$

and the convergence is uniform in $t > t_0$, due to monotonicity of of the integrand in $t > t_0$. On the other hand,

$$\left|\frac{2}{\pi}\int_B^\infty f_{n,t,x}(\lambda)d\lambda\right| \le I_n(t_0, B),$$

which converges to 0 as $B \to \infty$, uniformly in x and $t > t_0$ (by Remark 6.3). Hence, by a substitution $z = t\psi(\lambda^2)$,

$$\begin{aligned}\lim_{x\to 0^+}&\left(\sqrt{\psi(1/x^2)}\int_0^\infty f_{n,t,x}(\lambda)d\lambda\right)\\ &= \frac{1}{\Gamma(1+\alpha)}\int_0^\infty (\psi(\lambda^2))^{n-1/2}e^{-t\psi(\lambda^2)}\lambda\psi'(\lambda^2)d\lambda\\ &= \frac{1}{2\Gamma(1+\alpha)t^{n+1/2}}\int_0^\infty z^{n-1/2}e^{-z}dz = \frac{\Gamma(n+1/2)}{2\Gamma(1+\alpha)t^{n+1/2}}.\end{aligned}$$

Formula (1.13) follows now from (1.9). □

7. Examples

7.1. **Symmetric stable processes.** These processes, corresponding to $\Psi(\xi) = |\xi|^\alpha$ with $\alpha \in (0, 2]$, have already been studied in Example 6.1 in [28]. In this case $\vartheta_\lambda = (2-\alpha)\pi/8$, and Theorem 1.6 reads

$$(-1)^n \frac{d^n}{dt^n}\mathbf{P}_x(\tau_x > t) = \frac{\sqrt{2\alpha}}{\pi}\int_0^\infty \lambda^{n\alpha-1}e^{-t\lambda^\alpha}F(\lambda x)d\lambda, \qquad t, x > 0, \tag{7.1}$$

with (see Example 6.1 in [28])

$$\begin{aligned}F(\lambda x) = \sin(\lambda x + (2-\alpha)\pi/8) - \frac{\sqrt{2\alpha}\,\sin(\alpha\pi/2)}{2\pi}\int_0^\infty \frac{s^\alpha}{1+s^{2\alpha}-2s^\alpha\cos(\alpha\pi/2)}\\ \times\exp\left(\frac{1}{\pi}\int_0^\infty \frac{1}{1+\zeta^2}\log\frac{1-s^2\zeta^2}{1-s^\alpha\zeta^\alpha}\,d\zeta\right)e^{-\lambda s x}ds.\end{aligned}\tag{7.2}$$

Note that all above integrands are highly regular functions (for example, $\log((1-s^2\zeta^2)/(1-s^\alpha\zeta^\alpha))$ is a complete Bernstein function), and thus (7.2) is suitable for numerical integration. With a little effort, explicit upper bounds for numerical errors can also be computed. Together with (7.1), this gives *faithful* numerical bounds for $\mathbf{P}(\tau_x > t)$ and its derivatives in t. Plots of cumulative distribution function and density function of τ_x obtained using this method are given in Figure 1.

7.2. **Processes with power-type Lévy-Khintchine exponent.** In this example, we assume that the Lévy-Khintchine exponent $\Psi(\xi)$ has the form $\Psi(\xi) = \psi(\xi^2)$ for a complete Bernstein function ψ, and Ψ is regularly varying of *positive* order $\alpha_0 > 0$ at zero, and of *positive* order $\alpha_\infty > 0$ at $\pm\infty$ (see the first part of Table 1 for some examples). Clearly, in this case $\psi(\xi)$ is regularly varying of orders $\alpha_0/2$ and $\alpha_\infty/2$ at 0 and ∞, respectively. Hence, by Karamata's theory of regularly varying functions (see [8]), $-\xi\psi''(\xi)/\psi'(\xi)$ converges to $1-\alpha_0/2$ and $1-\alpha_\infty/2$ as $\xi \to 0^+$ and $\xi \to \infty$, respectively.

For simplicity, we assume in addition that the supremum in (1.7) is less than one. This condition is satisfied by all examples given in the first part of Table 1. Note, however, that the condition given in Remark 6.5(b) is automatically satisfied, so that our extra



(a)

(b)

(c)

Figure 1. Plots of $\mathbf{P}(\tau_x < t)$ (black) and $\frac{d}{dt}\mathbf{P}(\tau_x < t)$ (red) for the symmetric α-stable Lévy process, computed with three digits of accuracy, for (a) $\alpha = 0.5$; (b) $\alpha = 1.0$; (c) $\alpha = 1.5$. Calculations are based on Theorems 1.6 and 1.4 and the following numerical integration scheme.

Suppose that f is integrated over an interval I. First, using *analytical* methods, f is bounded above and below on any subinterval I' by a simpler function (e.g. a polynomial), which is integrated analytically. This yields lower and upper bounds for the integral of f over I'. A local adaptive strategy is used to divide I into sub-intervals I' so that the total error on all sub-intervals does not exceed given level. Interval arithmetic is used for nested integrals.

Plots prepared using `gnuplot` and a `C` program with 80-bit precision floating point numbers.

$\Psi(\xi)$	α_0	α_∞	X_t	restrictions
ξ^2	2	2	Brownian motion	
$\lvert\xi\rvert^\alpha$	α	α	α-stable	$\alpha \in (0,2]$
$c_1\lvert\xi\rvert^\alpha + c_2\lvert\xi\rvert^\beta$	α	β	(sum of two stables)	$\alpha, \beta \in (0,2]$, $\alpha < \beta$
$(\xi^2 + m^{2/\alpha})^{\alpha/2} - m$	2	α	relativistic α-stable	$\alpha \in (0,2)$, $m > 0$
$((\xi^2+1)^{\alpha/\beta} - 1)^{\beta/2}$	β	α	(subordinate relat. stable)	$\alpha, \beta \in (0,2]$, $\alpha < \beta$
$\log(1+\xi^2)$	2	0	variance gamma	
$\log(1+\lvert\xi\rvert^\alpha)$	α	0	geometric stable	$\alpha \in (0,2]$
$1/\log(1+1/\lvert\xi\rvert^\alpha)$	0	α	(not named)	$\alpha \in (0,2]$
$\log(1+1/\log(1+1/\xi^2))$	0	0	(not named)	
$\xi^2/(1+\xi^2)$	2	0	(compound Poisson)	

TABLE 1. Some Lévy-Khintchine exponents $\Psi(\xi)$, regularly varying both at 0 (of order α_0) and at $\pm\infty$ (of order α_∞). Names of corresponding subordinate Brownian motions are given in column X_t. First part of the table contains power-type functions Ψ, more singular examples are given in the other parts.

assumption can be easily dropped by referring to an improved version of Theorem 1.7, alluded to in Remark 6.5(b).

Theorem 1.7(b) yields the estimate

$$\frac{c_1(n)}{t^{n+1/2}\sqrt{\psi(1/x^2)}} \le (-1)^n \frac{d^n}{dt^n} \mathbf{P}(\tau_x > t) \le \frac{c_2(n)}{t^{n+1/2}\sqrt{\psi(1/x^2)}}$$

for $n \ge 0$, $x > 0$ and $t \ge c_3(n,\alpha)/\psi(1/x^2)$. Furthermore, by Theorem 1.8,

$$\begin{aligned} \lim_{x \to 0^+} \left(\sqrt{\psi(1/x^2)} \, \frac{d^n}{dt^n} \mathbf{P}(\tau_x > t) \right) &= \frac{(-1)^n \Gamma(n+1/2)}{\pi \Gamma(1+\alpha_\infty/2)} \, \frac{1}{t^{n+1/2}}, && t > 0,\ n \ge 0, \\ \lim_{t \to \infty} \left(t^{n+1/2} \, \frac{d^n}{dt^n} \mathbf{P}(\tau_x > t) \right) &= \frac{(-1)^n \Gamma(n+1/2)}{\pi} \, V(x), && x > 0,\ n \ge 0. \end{aligned}$$

7.3. **Slowly varying Lévy-Khintchine exponents.** When the Lévy-Khintchine exponent $\Psi(\xi)$ has the form $\Psi(\xi) = \psi(\xi^2)$ for a complete Bernstein function ψ, and Ψ is regularly varying of order α_0 at zero, and of order α_∞ at $\pm\infty$, but at least one of α_0, α_∞ is zero, estimates of the distribution of τ_x become more delicate. In this case Theorem 1.7 cannot be applied, and one needs to refer to either Corollary 6.2 (which is fairly straightforward, but typically yields sub-optimal results) or the technical Lemma 6.2. Note that only $n \ge 1$ need to be considered, as $n = 0$ was studied in general in [29].

Suppose that $|\psi''(\xi)|/\psi'(\xi) \le 1/\xi$ for all $\xi > 0$ (that is, $\varrho = 1$ in Theorem 1.7(b)); this condition is satisfied by all processes in the second part of Table 1. Then $\Theta = \sup_{\lambda > 0} \vartheta_\lambda \le \pi/4$ by Proposition 4.3. Furthermore, by integration, $\psi'(\xi_1)/\psi'(\xi_2) \le \xi_2/\xi_1$ when $0 < \xi_1 < \xi_2$, and therefore (cf. the proof of Theorem 1.7(b))

$$\begin{aligned} I_n(t, \lambda_0) &\le \frac{2}{\pi \sqrt{\lambda_0^2 \psi'(\lambda_0^2)}} \int_{\lambda_0}^\infty e^{-t\psi(\lambda^2)} (\psi(\lambda^2))^{n-1/2} (2\lambda\psi'(\lambda^2)) d\lambda \\ &= \frac{2}{\pi} \, \frac{\Gamma(n+1/2; t\psi(\lambda_0^2))}{t^{n+1/2}\sqrt{\lambda_0^2 \psi'(\lambda_0^2)}} \, , \end{aligned} \tag{7.3}$$

where $\Gamma(a;z)$ is the upper incomplete gamma function. However, $\lambda_0^2\psi'(\lambda_0^2)$ is no longer comparable with $\psi(\lambda_0^2)$. Nevertheless, we can combine (7.3) with Lemma 6.2, to find that

$$\begin{aligned}\frac{C_1(n,\pi/4)}{2}\,\frac{1}{t^{n+1/2}\sqrt{\psi(1/x^2)}} &\le (-1)^n \frac{d^n}{dt^n}\,\mathbf{P}(\tau_x > t) \\ &\le \left(C_1(n,\pi/4) + \frac{C_2(n,\pi/4)}{2}\right)\frac{1}{t^{n+1/2}\sqrt{\psi(1/x^2)}}\,,\end{aligned}$$

provided that $t\psi(1/x^2) \ge 1$ (so that the minimum in (6.4) is $1/(t^{n+1/2}(\psi(1/x^2))^{1/2})$) and

$$\Gamma(n+1/2; t\psi((\lambda_0(x))^2)) \le \frac{\pi C_1(n,\pi/4)}{4}\sqrt{\frac{(\lambda_0(x))^2\psi'((\lambda_0(x))^2)}{\psi(1/x^2)}}\,. \tag{7.4}$$

Here $C_1(n,\Theta)$, $C_2(n,\Theta)$ are the constants of Lemma 6.2, and $\lambda_0(x) = \pi/(8x)$.

Note that $\Gamma(n+1/2;z) \le c(n)e^{-z/2}$ for $z > 0$ for some $c(n)$. Furthermore, ψ' is a decreasing function. After some simplification (we omit the details), this gives the following sufficient condition for (7.4):

$$t\psi(1/x^2) \ge c'(n)\left(1 + \log\left(1 + \frac{\psi(1/x^2)}{(1/x^2)\psi'(1/x^2)}\right)\right)$$

for some $c'(n)$. For convenience, we state this as a separate result.

Proposition 7.1. *If $|\psi''(\xi)|/\psi'(\xi) \le 1/\xi$ for all $\xi > 0$, then there are positive constants $c_1(n)$, $c_2(n)$, $c_3(n)$ such that*

$$\frac{c_1(n)}{t^{n+1/2}\sqrt{\psi(1/x^2)}} \le (-1)^n\frac{d^n}{dt^n}\,\mathbf{P}(\tau_x > t) \le \frac{c_2(n)}{t^{n+1/2}\sqrt{\psi(1/x^2)}} \tag{7.5}$$

for $n \ge 0$, $x > 0$ and

$$t \ge \frac{c_3(n)}{\psi(1/x^2)}\left(1 + \log\left(1 + \frac{\psi(1/x^2)}{(1/x^2)\psi'(1/x^2)}\right)\right). \quad \square \tag{7.6}$$

For example, when $\Psi(\xi) = \log(1 + |\xi|^\alpha)$ ($\alpha \in (0,2]$), which corresponds to geometric stable processes, condition (7.6) reads

$$t \ge \frac{c(\alpha,n)}{\psi(1/x^2)}\,(1 + \log(1 + \log(1 + 1/x))).$$

In this case by Theorem 1.8 we also have

$$\begin{aligned}\lim_{x\to 0^+}\left(\sqrt{\log(1/x)}\,\frac{d^n}{dt^n}\,\mathbf{P}(\tau_x > t)\right) &= \frac{(-1)^n\Gamma(n+1/2)}{\pi\sqrt{\alpha/2}}\,\frac{1}{t^{n+1/2}}, && t > 0,\ n \ge 0, \\ \lim_{t\to\infty}\left(t^{n+1/2}\,\frac{d^n}{dt^n}\,\mathbf{P}(\tau_x > t)\right) &= \frac{(-1)^n\Gamma(n+1/2)}{\pi}\,V(x), && x > 0,\ n \ge 0.\end{aligned}$$

7.4. **Exponents with very slow growth.** Let $\tilde\psi(\xi) = \log(1 + \xi^\alpha)$ be the complete Bernstein function studied in the previous example, and let $\psi = \tilde\psi \circ \ldots \circ \tilde\psi$ be the N-fold composition of $\tilde\psi$ ($\alpha \in (0,2]$, $N \ge 2$). The function ψ is the Laplace exponent of the iterated geometric α-stable subordinator, and $\Psi(\xi) = \psi(\xi^2)$ is the Lévy-Khintchine exponent of the corresponding subordinate Brownian motion. It is easy to verify that (1.7) is satisfied. However, (1.8) holds only if $N = 2$ and $t \ge 1/\alpha$ (n arbitrary). Hence, for $N = 2$ asymptotic expansion (1.13) in Theorem 1.8(b) is valid only for $t > 1/\alpha$ (part (a) of the theorem holds for all $x > 0$). When $N \ge 3$, neither part of Theorem 1.8 applies.

Note that for any $N \ge 2$ and $\alpha \in (0,2]$, $\sup_{\xi>0}(\xi|\psi''(\xi)|/\psi'(\xi)) > 1$, so these examples do not fit into the framework of Theorem 1.7(b), or even that of Subsection 7.3. Some estimates of the derivatives of $\mathbf{P}(\tau_x > t)$ for $N = 2$ and $t > 1/\alpha$ can be obtained using Lemma 6.2, see e.g. Corollary 6.4.

7.5. **Compound Poisson process with Laplace distributed jumps.** Let $\Psi(\xi) = \xi^2/(1+\xi^2)$. Then the corresponding process X_t is the compound Poisson process with Laplace distributed (i.e. with density function $(1/2)e^{-|x|}$) jumps occuring at unit rate. It was proved in [29] that

$$\mathbf{P}(\tau_x > t) \approx \min\left(1, \sqrt{\frac{1+x^2}{t}}\right), \qquad x > 0,\, t > 0.$$

Note that (1.7) is not satisfied, so the main results of the present article cannot be used. Indeed, we have $\vartheta_\lambda = \arctan \lambda$ (see [28], Example 6.6), and so $\sup_{\lambda > 0} \vartheta_\lambda$ is indeed equal to $\pi/2$.

7.6. **Irregular example.** Estimates for ϑ_λ are critical for Theorems 1.7 and 1.8. We have $\vartheta_\lambda \in [0, \pi/2)$, and it is easy to construct examples for which $\lim_{\lambda\to\infty} \vartheta_\lambda$ is any number in $[0, \pi/2]$. On the other hand, it seems unlikely that $\liminf_{\lambda \to 0^+} \vartheta_\lambda > \pi/4$ is possible. Below we show a rather irregular example for which $\limsup_{\lambda\to 0^+} \vartheta_\lambda = \pi/2$ (but $\liminf_{\lambda\to 0^+} \vartheta_\lambda = 0$).

Consider the complete Bernstein function

$$\psi(\xi) = \sum_{k=1}^{\infty} p_k \, \frac{\xi}{a_k + \xi} \, ,$$

where, for example, $p_k = 1/k!$ and $a_k = 1/(k!)^2$. Fix $q > 0$. We consider $\lambda = (q a_n)^{1/2}$ and let $n \to \infty$. We have

$$\begin{aligned}
0 \le \psi(q a_n) \left(p_n \, \frac{q}{1+q}\right)^{-1} - 1 &= \sum_{k \ne n} \frac{p_k}{p_n} \, \frac{(1+q) a_n}{a_k + q a_n} \\
&\le \sum_{k=1}^{n-1} \frac{p_k}{p_n} \, \frac{(1+q) a_n}{a_k} + \sum_{k=n+1}^{\infty} \frac{p_k}{p_n} \, \frac{(1+q) a_n}{q a_n} \\
&= (1+q) \sum_{k=1}^{n-1} \frac{k!}{n!} + \frac{1+q}{q} \sum_{k=n+1}^{\infty} \frac{n!}{k!} \\
&\le \frac{1+q}{n} \sum_{k=1}^{n-1} \frac{1}{(n-k-1)!} + \frac{1+q}{q(n+1)} \sum_{k=n+1}^{\infty} \frac{1}{(k-n-1)!} \\
&\le \frac{(1+q)^2 e}{qn} = O(1/n).
\end{aligned}$$

Here and below the constant in the $O(\cdot)$ notation may depend on q. In a similar manner,

$$\begin{aligned}
0 \le \psi'(q a_n) \left(p_n \, \frac{1}{(1+q)^2 a_n}\right)^{-1} - 1 &= \sum_{k \ne n} \frac{p_k}{p_n} \, \frac{(1+q)^2 a_n a_k}{(a_k + q a_n)^2} \\
&\le \sum_{k=1}^{n-1} \frac{p_k}{p_n} \, \frac{(1+q)^2 a_n}{a_k} + \sum_{k=n+1}^{\infty} \frac{p_k}{p_n} \, \frac{(1+q)^2}{q} \\
&\le \frac{(1+q)^3 e}{qn} = O(1/n),
\end{aligned}$$

and

$$\begin{aligned} 0 \le |\psi''(qa_n)| \left(p_n \, \frac{2}{(1+q)^3 a_n^2} \right)^{-1} - 1 &= \sum_{k \ne n} \frac{p_k}{p_n} \, \frac{(1+q)^3 a_n^2 a_k}{(a_k + q a_n)^3} \\ &\le \sum_{k=1}^{n-1} \frac{p_k}{p_n} \, \frac{(1+q)^3 a_n}{a_k} + \sum_{k=n+1}^{\infty} \frac{p_k}{p_n} \, \frac{(1+q)^3}{q} \\ &\le \frac{(1+q)^4 e}{qn} = O(1/n). \end{aligned}$$

It follows that

$$\begin{aligned} \psi(qa_n) &= \left(p_n \, \frac{q}{1+q} \right) (1 + O(1/n)), \\ \psi'(qa_n) &= \left(p_n \, \frac{1}{(1+q)^2 a_n} \right) (1 + O(1/n)), \\ |\psi''(qa_n)| &= \left(p_n \, \frac{2}{(1+q)^3 a_n^2} \right) (1 + O(1/n)). \end{aligned}$$

With the notation of the proof of Proposition 4.5, for $\lambda^2 = qa_n$ we obtain

$$\begin{aligned} a_1^2 &= \frac{\psi(\lambda^2)}{\lambda^2 \psi'(\lambda^2)} = (1+q)(1 + O(1/n)), \\ a_2^2 &= \frac{\lambda^2 |\psi''(\lambda^2)|}{2\psi'(\lambda^2)} = \frac{q}{1+q} \, (1 + O(1/n)), \\ a^2 &= a_1^2 - 1 = q(1 + O(1/n)). \end{aligned}$$

By Proposition 4.5,

$$\limsup_{n \to \infty} \vartheta_{(qa_n)^{1/2}} \le \lim_{n \to \infty} \arctan \sqrt{q(1 + O(1/n))} = \arctan \sqrt{q},$$

and, in a similar manner,

$$\begin{aligned} \liminf_{n \to \infty} \vartheta_{(qa_n)^{1/2}} &\ge \frac{1}{\pi} \left(\left(\arcsin \sqrt{\frac{q}{1+q}} \right)^2 + (\arcsin 1)^2 - \left(\arcsin \sqrt{\frac{1}{1+q}} \right)^2 \right) \\ &= \frac{\pi}{4} - \frac{1}{\pi} \left(\arcsin \sqrt{\frac{1}{1+q}} + \arcsin \sqrt{\frac{q}{1+q}} \right) \left(\arcsin \sqrt{\frac{1}{1+q}} - \arcsin \sqrt{\frac{q}{1+q}} \right) \\ &= \frac{\pi}{4} - \frac{1}{\pi} \, \frac{\pi}{2} \, \arcsin \frac{1-q}{1+q} = \frac{1}{2} \, \arccos \frac{1-q}{1+q} = \frac{1}{2} \, \arctan \frac{2\sqrt{q}}{1-q} = \arctan \sqrt{q}. \end{aligned}$$

Hence, $\lim_{n \to \infty} \vartheta_{(qa_n)^{1/2}} = \arctan q^{1/2}$, and therefore any number in $[0, \pi/2]$ is a partial limit of ϑ_λ as $\lambda \to 0^+$.

REFERENCES

[1] M. Abramowitz, I. A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables.* Dover, New York, 9th edition, 1972.

[2] L. Alili, A. E. Kyprianou, *Some remarks on first passage of Lévy processes, the American put and pasting principles.* Ann. Appl. Probab. 15(3) (2005) 2062–2080.

[3] O. E. Barndorff-Nielsen, T. Mikosch, S. I. Resnick (Eds.), *Lévy Processes: Theory and Applications.* Birkhäuser, Boston, 2001.

[4] G. Baxter, M. D. Donsker, *On the distribution of the supremum functional for processes with stationary independent increments.* Trans. Amer. Math. Soc. 85 (1957) 73–87.

[5] V. Bernyk, R. C. Dalang, G. Peskir, *The law of the supremum of a stable Lévy process with no negative jumps.* Ann. Probab. 36(5) (2008) 1777–1789.

[6] J. Bertoin, *Lévy Processes.* Cambridge Univ. Press, Melbourne, New York, 1996.

[7] N. H. Bingham, *Maxima of sums of random variables and suprema of stable processes.* Z. Wahrscheinlichkeitstheorie Verw. Gebiete 26 (1973) 273–296.

[8] N. H. Bingham, C. M. Goldie, J. L. Teugels, *Regular Variation.* Cambridge University Press, Cambridge, 1987.

[9] K. Bogdan, T. Byczkowski, T. Kulczycki, M. Ryznar, R. Song, Z. Vondraček, *Potential Analysis of Stable Processes and its Extensions.* Lecture Notes in Mathematics 1980, Springer, 2009.

[10] D. A. Darling, *The maximum of sums of stable random variables.* Trans. Amer. Math. Soc. 83 (1956) 164–169.

[11] R. A. Doney, *On Wiener-Hopf factorisation and the distribution of extrema for certain stable processes.* Ann. Probab. 15(4) (1987) 1352–1362.

[12] R. A. Doney, *Fluctuation Theory for Lévy Processes.* Lecture Notes in Math. 1897, Springer, Berlin, 2007.

[13] R. A. Doney, V. Rivero, *Asymptotic behaviour of first passage time distributions for Lévy processes*/ Preprint, 2011, arXiv:1107.4415v1.

[14] R. A. Doney, M. S. Savov, *The asymptotic behavior of densities related to the supremum of a stable process.* Ann. Probab. 38(1) (2010) 316–326.

[15] P. Graczyk, T. Jakubowski, *On exit time of symmetric α-stable processes.* Stoch. Proc. Appl. 122 (2012) 31–41.

[16] P. Graczyk, T. Jakubowski, *On Wiener-Hopf factors of stable processes.* Ann. Inst. Henri Poincaré (B) 47(1) (2010) 9–19.

[17] P. E. Greenwood, A. A. Novikov, *One-sided boundary crossing for processes with independent increments.* Teor. Veroyatnost. i Primenen. 31(2) (1986) 266–277.

[18] F. Hubalek, A. Kuznetsov, *A convergent series representation for the density of the supremum of a stable process.* Elect. Comm. Probab. 16 (2011) 84–95.

[19] T. R. Hurd, A. Kuznetsov, *On the first passage time for Brownian motion subordinated by a Lévy process.* J. Appl. Prob. 46 (2009) 181–198.

[20] E. Katzav, M. Adda-Bedia, *The spectrum of the fractional Laplacian and First-Passage-Time statistics.* EPL 83, 30006 (2008).

[21] P. Kim, R. Song, Z. Vondraček, *Boundary Harnack principle for subordinate Brownian motions.* Stoch. Proc. Appl. 119 (2009) 1601–1631.

[22] P. Kim, R. Song, Z. Vondraček, *On the potential theory of one-dimensional subordinate Brownian motions with continuous components.* Potential Anal. 33 (2010) 153–173.

[23] P. Kim, R. Song, Z. Vondraček, *Two-sided Green function estimates for killed subordinate Brownian motions.* To appear in Proc. London Math. Soc., arXiv:1007.5455v2.

[24] P. Kim, R. Song, Z. Vondraček, *Potential theory of subordinate Brownian motions revisited.* To appear in a volume in Honor of Prof. Jiaan Yan, arXiv:1102.1369v2.

[25] T. Koren, J. Klafter, M. Magdziarz, *First passage times of Lévy flights coexisting with subdiffusion.* Phys. Rev. E 76 (2007) 031129.

[26] A. Kuznetsov, *Wiener-Hopf factorization and distribution of extrema for a family of Lévy processes.* Ann. Appl. Prob. 20(5) (2010) 1801–1830.

[27] A. Kuznetsov, *On extrema of stable processes.* Ann. Probab. 39(3) (2011) 1027–1060.

[28] M. Kwaśnicki, *Spectral analysis of subordinate Brownian motions in half-line.* Studia Math. 206(3) (2011) 211–271.

[29] M. Kwaśnicki, J. Małecki, M. Ryznar, *Suprema of Lévy processes.* To appear in Ann. Probab., arXiv:1103.0935v1.

[30] A. E. Kyprianou, *Introductory Lectures on Fluctuations of Lévy Processes with Applications.* Universitext, Springer-Verlag, Berlin, 2006.

[31] Z. A. Melzak, *Companion to Concrete Mathematics.* John Wiley & Sons, New York, 1973.

[32] R. Pyke, *The Supremum and Infimum of the Poisson Process.* Ann. Math. Stat. 30(2) (1959) 568–576.

[33] K. Sato, *Lévy Processes and Infinitely Divisible Distributions.* Cambridge Univ. Press, Cambridge, 1999.

[34] R. Schilling, R. Song, Z. Vondraček, *Bernstein Functions: Theory and Applications.* De Gruyter, Studies in Math. 37, Berlin, 2010.

[35] M. L. Silverstein, *Classification of coharmonic and coinvariant functions for a Lévy process.* Ann. Probab. 8(3) (1980) 539–575.

[36] V. M. Zolotarev, *The first-passage time of a level and the behaviour at infinity for a class of processes with independent increments.* Theor. Probab, Appl. 9 (1964) 653–664.

Mateusz Kwaśnicki, Jacek Małecki, Michał Ryznar, Institute of Mathematics and Computer Science, Wrocław University of Technology, ul. Wybrzeże Wyspiańskiego 27, 50-370 Wrocław, Poland
E-mail address: `mateusz.kwasnicki@pwr.wroc.pl`, `jacek.malecki@pwr.wroc.pl`, `michal.ryznar@pwr.wroc.pl`

Mateusz Kwaśnicki, Institute of Mathematics, Polish Academy of Sciences, ul. Śniadeckich 8, 00-976 Warszawa, Poland
E-mail address: `m.kwasnicki@impan.pl`

Jacek Małecki, LAREMA, Université d'Angers, 2 Bd Lavoisier, 49045 Angers cedex 1, France